



**Agromet Advisory Service Bulletin for Raigad District**  
(Issued jointly by IAAS, Dr. B.S. Konkan Krishi Vidyapeeth,  
& Regional India Meteorological Department, Mumbai)  
(02358) 282387



No. 36/2019

Date: 03/05/2019

Duration – 5 days

**Dr. Ashokkumar Chavan,**  
Head,  
Department of Agronomy  
9422373396

**Dr. Vijay More,**  
Nodal Officer,  
Department of Agronomy  
9422374001

**Dr. Shital Yadav,**  
Technical Officer,  
Department of Agronomy  
8379901160

Significant past weather for the preceding week (Period –27/04/2019 to 03/05/2019)							Weather Parameters	Weather forecast until 08.30 hrs of 08/05/2019				
27/04	28/04	29/04	30/04	01/05	02/05	03/05		04/05	05/05	06/05	07/05	08/05
-	-	-	-	-	-	-	Rainfall (mm)	0	0	0	0	0
-	-	-	-	-	-	-	Maximum temperature (°C)	35	35	34	34	34
-	-	-	-	-	-	-	Minimum temperature (°C)	24	25	25	25	26
-	-	-	-	-	-	-	Cloud cover (Octa)	5	7	2	0	1
-	-	-	-	-	-	-	Relative Humidity Max.(%)	87	86	86	83	84
-	-	-	-	-	-	-	Relative Humidity Min. (%)	37	42	44	41	42
-	-	-	-	-	-	-	Wind speed (Km/hr)	5	4	4	3	5
-	-	-	-	-	-	-	Wind direction	185	180	189	247	254
Rainfall (mm) in last week							Rainfall (mm) from 01/01/2019 to till dated	Total Rainfall (mm) in last year				
0.0							0.0	3558.3				

**Agro-met Advisory**  
Sky will be cloudy on 4<sup>th</sup> and 5<sup>th</sup> May, 2019.

According to NDVI, Agriculture vigour is moderate and according to SPI, mildly dry condition experienced in Raigad district.

Crop	Stage	Agro Advise
Summer rice	Maturity	<ul style="list-style-type: none"> <li>Drain out water from the field where rice crop are in maturity stage.</li> <li>Harvest the matured rice crop and thresh it immediately. Dry the threshed grains in sun for 2 to 3 days.</li> </ul>
Moong and Mataka	Flowering and pod stage	<ul style="list-style-type: none"> <li>Provide irrigation to moong and mataka crop where crop is in flowering and pod development stage.</li> </ul>
Mango	Fruiting (Egg to maturity stage)	<ul style="list-style-type: none"> <li>Harvest the mature fruits before 10 hours in the morning and after 16 hours in the evening with the help on Nutan mango harvester at 80 to 85% maturity immediately which will help to reduce spongy tissue. Keep the harvested fruits in shade to prevent from heat and spongy tissue.</li> <li>To prevent incidence of post harvest diseases on fruits, place the fruits in hot water of 52°C for 10 minutes and then keep for ripening. Use C.F.B. (corrugated fiber box) for packing developed by B. S. Konkan Krishi Vidyapeeth, Dapoli. Transport of harvested fruits should be done preferably during night hours. Do not spray any insecticides/fungicides 8 days before harvesting of mango fruits.</li> <li>Due to forecast of cloudy weather, there is possibility of incidence of anthracnose disease and fruit fly incidence on mango. For control of anthracnose spray Carbendazim 12% + Mancozeb 63% @ 10 g per 10 liter of water. For control of fruit fly incidence, install 'Rakshak fruit fly trap' developed by University @ 4 traps per hectare. Collect and destroy fallen fruits and keep orchard clean.</li> <li>Clean mango tree by removing loranthus and all dried diseased infected branches and apply bordopaste on cut portion of tree. Maintain good sanitation in orchard.</li> </ul>
Cashewnut	-	<ul style="list-style-type: none"> <li>Clean cashewnut tree by removing all dried infected branches. Maintain good sanitation in orchard. Apply bordopaste on cut portion of tree.</li> <li>If the incidence of Cashew Stem and Root Borer (CSRB) is noticed in Cashew, remove the grubs from the holes with the help of 15 mm chisel and apply Chloropyrifos 20 EC @ 5 ml per liter of water on the stem (swabbing) and pour remaining solution around the stem near to the roots.</li> </ul>
Arecanut	-	<ul style="list-style-type: none"> <li>Clean the arecanut plant by removing dry leaves and diseased, infected inflorescence and spray 1% bordomixture to control 'Koleroga' disease. Repeat the spray 3-4 times at one month interval.</li> </ul>
Vegetables/ Fruit crop nursery	-	<ul style="list-style-type: none"> <li>Due to increase in rate of evaporation, provide irrigation regularly to fruit crop nursery, new planted fruit crops and vegetable crops.</li> </ul>
Milch animal /goat/poultry	-	<ul style="list-style-type: none"> <li>Provide clean, hygienic and plenty amount of drinking water to farm animals and poultry birds. To reduce the stress of heat in farm animals, provide roughages by mixing with solution of 1% gaggery and 0.5% salt separately.</li> <li>There is forecast for increase in temperature, hence protect animals and poultry birds from heat by covering roof of the shed with insulating materials such as paddy straw, dry coconut leaves and make arrangement for sprinkle cold water on the roof of shed during afternoon time. Use wet gunny bags as side curtains to protect animals and poultry birds from direct hot winds.</li> <li>To protect animals from heat, sprinkle cold water on animals during the afternoon, it will help to maintain the body temperature.</li> <li>In poultry shed, increase the water pot and provide adequate and clean water for drinking. Also, feed should be given in the morning or evening hours.</li> <li>Vaccination against foot and mouth disease in farm animals under supervision of veterinary officers is advocated.</li> </ul>

This Agro Advisory Bulletin (AAB) is prepared and published with the consultation and recommendation of SMS committees of "Gramin Krishi Mausam Sewa (GKMS)" Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli.

For more information contact nearby SAU research station or Agriculture officers of Agriculture Department, Maharashtra state.