



**Agromet Advisory Service Bulletin for Ratnagiri District**  
(Issued jointly by IAAS, Dr. B.S. Konkan Krishi Vidyapeeth,  
& Regional India Meteorological Department, Mumbai)  
(02358) 282387



No. 11/2019

Date: 05/02/2019

Duration – 5 days

**Dr. Subhash Chavan,**  
Head,  
Department of Agronomy  
9422431067

**Dr. Vijay More,**  
Nodal Officer,  
Department of Agronomy  
9422374001

**Dr. Shital Yadav,**  
Technical Officer,  
Department of Agronomy  
8379901160

Significant past weather for the preceding week (Period –30/01/2019 to 05/02/2019)							Weather Parameters	Weather forecast until 08.30 hrs of 10/02/2019				
30/01	31/01	01/02	02/02	03/02	04/02	05/02		06/02	07/02	08/02	09/02	10/02
0.0	0.0	0.0	0.0	0.0	0.0	0.0	Rainfall (mm)	0	0	0	0	0
31.0	31.6	31.2	31.2	32.2	32.4	34.2	Maximum temperature (°C)	34	31	30	30	31
12.0	11.8	13.0	13.0	14.5	14.0	14.5	Minimum temperature (°C)	20	18	17	15	15
0	0	0	0	0	0	0	Cloud cover (Octa)	0	1	0	0	0
93	95	91	90	91	89	87	Relative Humidity Max. (%)	68	65	76	60	49
60	61	66	65	61	61	66	Relative Humidity Min. (%)	46	52	49	40	30
3.8	3.1	4.2	3.3	3.1	3.3	3.5	Wind speed (Km/hr)	12	18	22	21	8
calm	calm	calm	calm	calm	calm	calm	Wind direction	38	316	321	17	71
Rainfall (mm) in last week							Rainfall (mm) from 01/01/2019 to till dated	Total Rainfall (mm) in last year				
0.0							0.0	3071.8				

**Agro-met Advisory**

There will be normal weather condition from 6<sup>th</sup> to 10<sup>th</sup> February, 2019. Wind velocity range from 8 to 22 Km/h from 6<sup>th</sup> to 10<sup>th</sup> February, 2019.

Crop	Stage	Agro Advise
Groundnut	Flowering	<ul style="list-style-type: none"> <li>Due to increasing rate of evaporation irrigate the groundnut crop at an interval of 8-10 days after sowing. Rolling of empty drum 45-60 days after sowing for better penetration of pegs &amp; pod setting is advocated.</li> </ul>
Pulse crop	Pod stage	<ul style="list-style-type: none"> <li>Harvest mature pulse crop early in morning or late in evening and dry it for 3 to 4 days in sunlight. Stored dried grain in proper manner.</li> </ul>
Mango	Flowering and Fruiting	<ul style="list-style-type: none"> <li>Provide support to newly planted mango graft, as there is forecast of high wind speed.</li> <li>Due to forecast of cloudy weather condition, there is possibility for incidence of hoppers and anthracnose disease on mango tree. Therefore it is advise to spray Thiomethoxam 25%WG @ 1gm for control the incidence of mango hoppers + Hexaconazole 5% @ 5 ml or wettable Sulphur 80% @ 20 gm for control of powdery mildew + Carbendazim 12% +Mancozeb 63% @ 10 gm for control of anthracnose disease per 10 liter of water.</li> <li>Provide irrigation to newly planted mango orchard and use straw mulch to reduce evaporation.</li> </ul>
Cashewnut	Fruiting	<ul style="list-style-type: none"> <li>Provide support to newly planted cashew graft, as there is forecast of high wind speed.</li> <li>There is possibility of incidence of tea mosquito bug and thrips during fruit bearing stage of cashewnut, if incidence is noticed spray Lambda cyhalothrin 5%EC @6 ml per 10 liter of water. (insecticide is not under label claim)</li> <li>Provide irrigation to newly planted cashewnut orchard and use straw mulch to reduce evaporation.</li> </ul>
Coconut	-	<ul style="list-style-type: none"> <li>For above five year old coconut palm apply 3<sup>rd</sup> dose of 750 gm urea and 667 gm muriate of potash in circular basin of 1.5 to 1.8 meter from the base of palm, incorporate in soil and irrigate.</li> </ul>
Arecanut	-	<ul style="list-style-type: none"> <li>Provide irrigation to arecanut orchard at 7-8 days interval.</li> <li>If incidence of red ant is noticed on Arecanut palm, dusting the stem with Deltamethrin 25%WP.</li> </ul>
Spice crop	Harvesting	<ul style="list-style-type: none"> <li>Cary out harvesting of black paper by picking the whole spikes by hand, when few berries in the spike start turning yellow or red. The berries are separated and dipped in hot water (80°C) for one minute and sun dried for 3 to 4 days.</li> </ul>
Vegetables/ Fruit crop nursery	Fruiting	<ul style="list-style-type: none"> <li>Follow earthing up operation in Okra, Brinjal, Tomato, Chilli etc. vegetable crops. Also provide support to crop as there is forecast of high wind speed.</li> <li>Carryout harvesting of the vegetable those are in proper stage of harvesting.</li> <li>Provide irrigation to fruit crop nursery, vegetable crops.</li> </ul>
Milch animal /goat/poultry	-	<ul style="list-style-type: none"> <li>Protect the animals /poultry birds from low temperature by providing curtains and electric bulbs as per the need.</li> <li>Provide clean and hygienic drinking water to farm animals.</li> <li>Medication against ticks in farm animal under the guidance of veterinary officer is advocated.</li> </ul>

This Agro Advisory Bulletin (AAB) is prepared and published with the consultation and recommendation of SMS committees of  
“Gramin Krishi Mausam Sewa (GKMS)” Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli.

For more information contact nearby SAU research station or Agriculture officers of Agriculture Department, Maharashtra state.