



Agromet Advisory Service Bulletin for Ratnagiri District
(Issued jointly by IAAS, Dr. B.S. Konkan Krishi Vidyapeeth,
& Regional India Meteorological Department, Mumbai)
(02358) 282387



No. 30/2019

Date: 12/04/2019

Duration – 5 days

Dr. Ashokkumar Chavan,
Head,
Department of Agronomy
9422373396

Dr. Vijay More,
Nodal Officer,
Department of Agronomy
9422374001

Dr. Shital Yadav,
Technical Officer,
Department of Agronomy
8379901160

Significant past weather for the preceding week (Period –06/04/2019 to 12/04/2019)							Weather Parameters	Weather forecast until 08.30 hrs of 17/04/2019				
06/04	07/04	08/04	09/04	10/04	11/04	12/04		13/04	14/04	15/04	16/04	17/04
0.0	0.0	0.0	0.0	0.0	0.0	0.0	Rainfall (mm)	0	5	0	0	3
32.0	33.4	33.0	35.2	32.4	33.5	32.8	Maximum temperature (°C)	34	35	36	35	35
20.5	18.0	22.0	21.0	20.0	21.5	23.5	Minimum temperature (°C)	25	27	26	26	25
0	0	0	0	0	0	0	Cloud cover (Octa)	0	1	1	2	1
89	91	89	91	91	89	90	Relative Humidity Max. (%)	69	58	66	81	75
57	52	43	54	48	50	50	Relative Humidity Min. (%)	28	24	24	40	39
4.9	5.5	5.2	6.0	5.1	6.9	5.6	Wind speed (Km/hr)	3	5	5	3	3
Calm	Calm	Calm	Calm	Calm	Calm	Calm	Wind direction	120	181	122	134	126
Rainfall (mm) in last week						Rainfall (mm) from 01/01/2019 to till dated			Total Rainfall (mm) in last year			
0.0						0.0			3071.8			

Agro-met Advisory

It is possibility rainfall of 14th and 17th April, 2019. There will be increase in maximum and minimum temperature from 13th to 17th April 2019.
According to NDVI, Agriculture vigour is moderate and according to SPI, mildly wet condition experienced in Ratnagiri district.

Crop	Stage	Agro Advise
Groundnut	Pod development	<ul style="list-style-type: none"> Provide irrigation to groundnut when crop is in pod filling stage.
Mango	Fruiting (Egg to maturity stage)	<ul style="list-style-type: none"> Harvest the mature fruits if any before 10 hours in the morning and after 16 hours in the evening with the help on Nutan mango harvester at 80 to 85% maturity. Keep the harvested fruits in shade to prevent spongy tissue disease and from heat. To prevent incidence of post harvest diseases on fruits, place the fruits in hot water of 52°C for 10 minutes and then keep for ripening. Use C.F.B. (corrugated fiber box) for packing developed by B. S. Konkan Krishi Vidyapeeth, Dapoli. Transport of harvested fruits should be done preferably during night hours. Do not spray any insecticides/fungicides 8 days before harvesting of mango fruits. There is forecast of light rainfall on 14th and 17th April, for protection of mango fruits from incidence of anthracnose and powdery mildew disease, spray Carbendazim 12% + Mancozeb 63% @ 10 g per 10 liter of water for control of anthracnose + Hexaconazole 5% @ 5 ml or wettable Sulphur 80% @ 20 gm per 10 liter of water for control of powdery mildew disease. Further for control of fruit fly incidence, install 'Rakshak fruit fly trap' developed by University @ 4 traps per hectare. Collect and destroy fallen fruits and keep orchard clean. To protect the mango fruits from raindrop, cover the fruits with paper bag. Provide irrigation to newly planted mango orchard @ 30 liters of water twice in week (1 years old), twice in 15 days interval (2 years old) and twice in month (3 years old). New growth below graft union should be removed regularly. For new plantation of mango, clean the field and dig the pits of size 1 X 1 X 1 m at a spacing of 10 X 10 m (5 X 5 m for high density planting). Re-filled the pits with mixture of soil, 3 to 4 basket of well decomposed FYM and 3 kg of single super phosphate. <p>Note: partially decomposed FYM will act as attractant for termite.</p>
Cashewnut	Fruiting and maturity	<ul style="list-style-type: none"> If nuts are ready for harvesting, harvest the mature nuts and dry in the sun for 7 to 8 days to bring down the moisture content. Prepare different processed product from cashew apple as per recommended by Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli. Clean cashewnut orchard by removing all diseased and dried branches to maintain good sanitation. Provide irrigation to newly planted cashew orchard @ 15 liters of water once in week. New growth below graft union should be removed regularly. For new plantation of cashewnut, clean the field, dig the pits of size 0.6 X 0.6 X 0.6 m at a spacing of 7.5 X 7.5 m or 8 X 8 m. Re-filled the pits with mixture of soil, 1½ to 2 basket of well decomposed FYM and ½ kg of single super phosphate. <p>Note: partially decomposed FYM will act as attractant for termite.</p>
Vegetables/ Fruit crop nursery	flowering	<ul style="list-style-type: none"> There is possibility of incidence of shoot and fruit borer on okra, if incidence is noticed spray cypermethrin 25%EC @ 3ml or Lambda cyhalothrin 5%EC @ 6ml per 10 liter of water. Due to increase in rate of evaporation, provide irrigation regularly to fruit crop nursery, new planted fruit crops and vegetable crops.
Milch animal /goat/poultry	-	<ul style="list-style-type: none"> Provide clean, hygienic and plenty amount of drinking water to farm animals and poultry birds. There is forecast for increase in temperature, hence protect animals and poultry birds from heat by covering roof of the shed with insulating materials such as paddy straw, dry coconut leaves and make arrangement for sprinkle cold water on the roof of shed during afternoon time. Use wet gunny bags as side curtains to protect animals and poultry birds from direct hot winds. To protect the farm animals from attack of ticks, apply 2% methyl parathion powder on to body.

This Agro Advisory Bulletin (AAB) is prepared and published with the consultation and recommendation of SMS committees of "Gramin Krishi Mausam Sewa (GKMS)" Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli.
For more information contact nearby SAU research station or Agriculture officers of Agriculture Department, Maharashtra state.