



Agromet Advisory Service Bulletin for Ratnagiri District
(Issued jointly by IAAS, Dr. B.S. Konkan Krishi Vidyapeeth,
& Regional India Meteorological Department, Mumbai)
(02358) 282387



No. 43/2019

Date: 28/05/2019

Duration – 5 days

Dr. Ashokkumar Chavan,
Head,
Department of Agronomy
9422373396

Dr. Vijay More,
Nodal Officer,
Department of Agronomy
9422374001

Dr. Shital Yadav,
Technical Officer,
Department of Agronomy
8379901160

Significant past weather for the preceding week (Period –22/05/2019 to 28/05/2019)							Weather Parameters	Weather forecast until 08.30 hrs of 02/06/2019				
22/05	23/05	24/05	25/05	26/05	27/05	28/05		29/05	30/05	31/05	01/06	02/06
0.0	0.0	0.0	0.0	0.0	0.0	0.0	Rainfall (mm)	0	0	0	0	0
33.2	34.2	34.0	34.4	34.0	33.8	33.2	Maximum temperature (°C)	34	35	35	35	34
24.0	23.0	24.3	25.0	22.5	22.5	24.5	Minimum temperature (°C)	27	27	27	28	28
2	2	4	4	4	2	2	Cloud cover (Octa)	2	7	6	3	4
80	80	92	87	83	80	78	Relative Humidity Max. (%)	90	92	91	89	88
60	65	58	55	50	59	58	Relative Humidity Min. (%)	42	43	45	45	49
6.4	6.5	7.0	5.9	6.8	5.9	6.5	Wind speed (Km/hr)	7	7	6	6	6
Calm	Calm	Calm	Calm	Calm	Calm	Calm	Wind direction	276	256	251	271	289
Rainfall (mm) in last week						Rainfall (mm) from 01/01/2019 to till dated			Total Rainfall (mm) in last year			
0.0						0.0			3071.8			

Agro-met Advisory

Sky will be cloudy from 29th May to 2nd June, 2019.

According to NDVI, Agriculture vigour is moderate and according to SPI, moderately dry condition experienced in Ratnagiri district.

Crop	Stage	Agro Advise
Kharif rice	Nursery preparation	<ul style="list-style-type: none"> For preparation of rice nursery, plough the area add FYM@100 kg per guntha area and bring the soil to fine tilth. Prepare raised bed of 120 cm. breadth at bottom and 90 cm. on top along the slope of land. Convenient length of raised bed should be kept according to the land slope. Store sufficient quantity of improved seed and required fertilizers during kharif season.
Finger millet	Nursery preparation	<ul style="list-style-type: none"> For preparation of finger millet nursery, plough the area add FYM@250 kg per guntha area and bring the soil to fine tilth. Prepare raised bed of 120 cm. breadth at bottom and 90 cm. on top along the slope of land. Convenient length of raised bed should be kept according to the land slope.
Mango	Fruiting	<ul style="list-style-type: none"> Harvest the mature fruits before 10 hours in the morning and after 16 hours in the evening with the help on Nutan mango harvester at 80 to 85% maturity immediately which will help to reduce spongy tissue. Keep the harvested fruits in shade to prevent from heat and spongy tissue. To prevent incidence of post harvest diseases on fruits, place the fruits in hot water of 52°C for 10 minutes and then keep for ripening. Use C.F.B. (corrugated fiber box) for packing developed by B. S. Konkan Krishi Vidyapeeth, Dapoli. Transport of harvested fruits should be done preferably during night hours. Do not spray any insecticides/fungicides 8 days before harvesting of mango fruits. For control of fruit fly incidence, install 'Rakshak fruit fly trap' developed by University @ 4 traps per hectare. Collect and destroy fallen fruits and keep orchard clean.
Coconut	-	<ul style="list-style-type: none"> There is possibility of incidence of bud rot fungal disease on young coconut palm during rainy season. To prevent the incidence of bud rot disease, spray 1% bordomixture onto the crown region of palm.
Vegetables/ Fruit crop nursery	-	<ul style="list-style-type: none"> For nursery of brinjal, chilli and tomato vegetables crops, prepare raised bed of 3m length x 1 m breadth x 15cm height. Apply 5 kg FYM, 35gm urea, 100gm single super phosphate and 25gm muriate of potash per sq. m. Sow the vegetable seed if irrigation facility is available. Treat the seed with thiram fungicide @ 3 gm/kg of seed before sowing. To protect the seedling from incidence of wilt disease, drenching of 1% bordomixture 3 to 4 days before sowing on nursery bed is suggested. Provide sufficient irrigation to fruit crop nursery also provide shed to nursery seedlings.
Milch animal /goat/poultry	-	<ul style="list-style-type: none"> Provide clean, hygienic and plenty amount of drinking water to farm animals and poultry birds. To reduce the stress of heat in farm animals, provide roughages by mixing with solution of 1% jaggery and 0.5% salt separately. There is forecast for increase in temperature, hence protect animals and poultry birds from heat by covering roof of the shed with insulating materials such as paddy straw, dry coconut leaves and make arrangement for sprinkle cold water on the roof of shed during afternoon time. Use wet gunny bags as side curtains to protect animals and poultry birds from direct hot winds. To protect animals from heat, sprinkle cold water on animals during the afternoon, it will help to maintain the body temperature. Due to increase in temperature, avoid grazing of farm animals during afternoon hours. In poultry shed, increase the water pot and provide adequate and clean water for drinking. Also, feed should be given in the morning or evening hours.

This Agro Advisory Bulletin (AAB) is prepared and published with the consultation and recommendation of SMS committees of "Gramin Krishi Mausam Sewa (GKMS)" Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli.

For more information contact nearby SAU research station or Agriculture officers of Agriculture Department, Maharashtra state.