

## Agromet Advisory Service Bulletin for Palghar District (Issued jointly by IAAS, Dr. B.S. Konkan Krishi Vidyapeeth, & Regional India Meteorological Department, Mumbai) (02358) 282387



No. 80/2019 Date: 04/10/2019 Duration – 5 days

**Dr. Prashant Bodake,**Head,
Department of Agronomy
9420413255

**Dr. Vijay More,**Nodal Officer,
Department of Agronomy
9422374001

**Dr. Shital Yadav,**Technical Officer,
Department of Agronomy
8379901160

Si	gnificant (Per	•		the prec o 04/10/2	0	veek	Weather Parameters	Weather forecast until 08.30 hrs of 09/10/2019				
28/09	29/09	30/09	01/10	02/10	03/10	04/10		05/10	06/10	07/10	08/10	09/10
4.2	0.0	22.2	7.6	0.0	0.0	0.0	Rainfall (mm)	0	0	16	15	13
29.0	31.0	32.4	30.6	31.0	31.0	32.4	Maximum temperature (°C)	32	32	33	34	34
24.8	27.4	26.4	26.0	26.0	26.0	26.4	Minimum temperature (°C)	24	25	25	26	26
1	1	2	0	0	0	0	Cloud cover (Octa)	4	4	6	6	8
95	85	92	92	92	84	80	Relative Humidity Max. (%)	83	83	80	78	76
74	72	83	77	77	69	-	Relative Humidity Min. (%)	68	66	66	63	61
4.6	5.3	4.1	3.0	2.4	1.9	2.0	Wind speed (Km/hr)	11.3	11.4	12.5	11.1	8.3
NNW	NNW	N	NNW	W	ESE	ESE	Wind direction	NNE	N	N	SSW	ENE
Rainfall (mm) in last week							Rainfall (mm) from 01/01/2019 to till dated	Total Rainfall (mm) in last		ast year		
34.0							4156.4	2359.6				

Agro-met Advisory

There is possibility of medium rainfall from 7<sup>th</sup> to 9<sup>th</sup> October, 2019 also possibility of increase in maximum temperature and sky will be cloudy.

Extended range rainfall forecast for Konkan division for the period from 4<sup>th</sup> to 10<sup>th</sup> October, 2019 is normal. According to NDVI, Agriculture vigour is moderate and according to SPI, severely wet condition experienced in Palghar district.

	_	district.
Crop	Stage	Agro Advise
Kharif rice	Flowering to grain filling stage	<ul> <li>Maintain 5 cm level of water in rice field where mid- late rice varieties in grain filling stage and late rice varieties is in flowering stage.</li> <li>Drain out water from the field where early rice varieties is in maturity stage.</li> <li>There is forecast of moderate rain on 7<sup>th</sup>, 8<sup>th</sup> and 9<sup>th</sup> October, hence by observing rainfall situation harvest matured early rice varieties in morning hours only with 'Vaibhav sickle' near to ground level and follow immediate threshing, the threshed grains should be sun dry for 2 to 3 days before storage.</li> <li>Due to forecast of increase in temperature and humidity, there is possibility of incidence of army worm in early rice varieties, hence by observing rainfall situation harvest matured early rice varieties.</li> </ul>
Mango	Vegetative	<ul> <li>Due to high humidity condition in past created favourable condition for growth of fungus in dense trees. Due to forecast of increase in temperature, the fungal growth will increase and result in dieback of branches in mango. For control of disease, collect and destroy all diseased infected leaves, branches and apply bordopaste to cut portion of branches. After cleaning of orchard, spray Metalaxyl 8% + mancozeb 64% combination fungicide@20 g per 10 liter of water.</li> <li>Due to increase in temperature and humidity there is possibility of incidence of hoppers, midge fly and shoot borer on vegetative flush of mango to protect the flush of mango, spray Lambda cyhalothrin 5%EC @ 6ml or Quinalphos 25%EC@25 ml per 10 liter of water.</li> </ul>
Cashewnut	Vegetative	<ul> <li>Due to high humidity condition in past created favourable condition for growth of fungus in dense trees. Due to forecast of increase in temperature, the fungal growth will increase and result in dieback of branches in cashew. For control of disease, collect and destroy all diseased infected leaves, branches and apply bordopaste to cut portion of branches. After cleaning of orchard, spray Metalaxyl 8% + mancozeb 64% combination fungicide@20 g per 10 liter of water.</li> <li>Due to increase in temperature and humidity there is possibility of incidence of tea mosquito bug and thrips on vegetative flush of cashewnut to protect the flush of cashew, spray Monocrotophos 36%SL @ 15 ml or Lambda cyhalothrin 5% EC @ 6 ml per 10 liter of water.</li> </ul>
Coconut	Fruiting	• For control of eriophyid mite on coconut, neem-based insecticide neemazal 5% @ 7.5 ml be mixed in equal quantity of water apply through root feeding during month of October-November. Harvesting is avoided at least 45 days after treatment. In addition to this spray neem-based insecticide (nemazal) 1% @4 ml per liter of water on bunch of nuts. Collect and destroy all infected inflorescence and nuts before spraying.
Sapota	Fruiting	• Follow the weeding operation in sapota orchard, also remove all dried, unproductive and

		disease infected branches, apply bordopaste on cut portion. After cleaning of orchard, spray 1% bordo mixture on to tree.For control of seed borer incidence on sapota, after cleaning of orchard, spray Profenophos 50% EC @ 15 ml or Indoxacarb 14.5 SC@ 5 ml or Novaluron 10%EC@ 5 ml or Deltamethrin 2.8 EC @ 10 ml insecticide per 10 liter of water. Repeat the
		spray at an interval of a month. Harvest all matured fruits before spraying.
Vegetables crop nursery	Sowing	• For raising of vegetable nursery for rabi season, cultivate the land when it is at field capacity condition and then prepare raised bed of 3m length x 1 m breadth x 15cm height. Apply 5 kg FYM, 35gm urea, 100gm single super phosphate and 25gm muriate of potash per sq. m. For protection of seedlings from wilt disease, drench nursery bed with 1% bordomixture before 3-4 days of sowing. Before sowing, treat the seed with Thiram or Captan fungicide 3 grams per kg of seed.
Milch animals/Goat		• Provide clean and hygienic water to farm animals also maintains good aeration in farm shed.

This Agro Advisory Bulletin (AAB) is prepared and published with the consultation and recommendation of SMS committees of "Gramin Krishi Mausam Sewa (GKMS)" Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli. For more information contact nearby SAU research station or Agriculture officers of Agriculture Department, Maharashtra state.