



Agromet Advisory Service Bulletin for Ratnagiri District
(Issued jointly by GKMS, Dr. B.S. Konkan Krishi Vidyapeeth,
& Regional India Meteorological Department, Mumbai)
(02358) 282387



No. 19/2020

Date: 06/03/2020

Duration – 5 days

Dr. Prashant Bodake,
Head,
Department of Agronomy
9420413255

Dr. Vijay More,
Nodal Officer,
Department of Agronomy
9422374001

Dr. Shital Yadav,
Technical Officer,
Department of Agronomy
8379901160

| Significant past weather for the preceding week (Period –29/02/2020 to 06/03/2020) | | | | | | | Weather Parameters | Weather forecast until 08.30 hrs of 11/03/2020 | | | | |
|---|-------|-------|-------|-------|-------|-------|--|---|-------|-------|-------|-------|
| 29/02 | 01/03 | 02/03 | 03/03 | 04/03 | 05/03 | 06/03 | | 07/03 | 08/03 | 09/03 | 10/03 | 11/03 |
| 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | Rainfall (mm) | 0 | 0 | 0 | 0 | 0 |
| 35.0 | 34.4 | 33.0 | 32.5 | 30.8 | 29.8 | 29.0 | Maximum temperature (°C) | 31 | 31 | 30 | 29 | 29 |
| 15.0 | 15.5 | 15.0 | 14.0 | 14.5 | 14.5 | 17.0 | Minimum temperature (°C) | 21 | 21 | 20 | 19 | 19 |
| 0 | 0 | 0 | 0 | 0 | 0 | 4 | Cloud cover (Octa) | 0 | 1 | 1 | 1 | 2 |
| 85 | 90 | 89 | 90 | 91 | 93 | 92 | Relative Humidity Max. (%) | 89 | 89 | 80 | 91 | 90 |
| 48 | 54 | 48 | 47 | 50 | 59 | - | Relative Humidity Min. (%) | 40 | 28 | 27 | 35 | 41 |
| 3.7 | 3.2 | 3.4 | 3.8 | 3.6 | 3.8 | 3.6 | Wind speed (Km/hr) | 5 | 4 | 6 | 4 | 4 |
| Calm | Calm | Calm | Calm | Calm | Calm | Calm | Wind direction | E | ESE | SSW | E | ESE |
| Rainfall (mm) in last week | | | | | | | Rainfall (mm) from 01/01/2020 to till dated | Total Rainfall (mm) in last year | | | | |
| 0.0 | | | | | | | 0.0 | 5130.9 | | | | |

Agro-met Advisory
Sky remain clear from 7th to 11th March, 2020.

| Crop | Stage | Agro Advise |
|-------------|-----------------------|--|
| Lablab bean | Harvesting | <ul style="list-style-type: none"> Harvest mature lablab bean pod and dry it for 4 to 5 days in sunlight and then follow threshing or harvest the pods along with plant and dry for 3 to 4 days in sunlight. After drying follow threshing of pods. Stored dried grain in proper manner. |
| Mango | Flowering to fruiting | <ul style="list-style-type: none"> To protect the flower bud of mango from hoppers, thrips and powdery mildew diseases, as per blossom protection schedule for mango crop, take a third spray of Imidacloprid 17.8% SL @ 3 ml per 10 liter of water before the flower opening (15 days after 2nd spray) to avoid the adverse effect on pollinators. Also add Hexaconazole 5% @ 5 ml or wettable Sulphur 80% @ 20 gm per 10 liter in water for control of powdery mildew disease. Note: Avoid spraying during flowering to fruit setting period for effective pollination. If it is not possible to postpone the spraying till fruit set due to heavy incidence of insect and pest, then avoid spraying during morning hours (10.00 am to 12.00 pm) which is active period of pollinators for pollination. To protect the pea size fruits of mango from hoppers, thrips and powdery mildew disease, as per blossom protection schedule for mango crop, take a fourth spray of Thiomethoxam 25%WG @ 1 gm per 10 liter of water (15 days after 3rd spray) Also add Hexaconazole 5% @ 5 ml or wettable Sulphur 80% @ 20 gm per 10 liter in water for control of powdery mildew. To minimize the pre-mature fruit drop of mango, apply 150 to 200 liter of water per tree after fruit setting (pea size) at 15 days interval for 3 to 4 times also use straw mulch to reduce evaporation losses. To improve production and quality of mango fruits, spray 1% Potassium nitrate at pea, marble and egg fruit stages. The pre-harvest bagging with newspaper bag of size 25 X 20 cm at marble to egg stage as per recommendation of D.B.S.K.K.V. helps to reduce the fruit drop, increases the fruit weight, pulp weight, produce spongy tissue free fruit, controls attack of fruit fly on fruits and produces spotless fruits of mango. Spraying of 55% cow urine at pea size fruits of mango 3 to 6 sprays at weekly interval as per availability is suggested for increasing yield. |
| Cashewnut | Flowering to fruiting | <ul style="list-style-type: none"> There is possibility of incidence of tea mosquito bug and thrips on new vegetative flush of cashewnut, to protect the flush of cashew spray Monocrotophos 36%SL @ 15 ml or Lambda cyhalothrin 5% EC @ 6 ml per 10 liter of water. To improve fruit setting and production of cashewnut, spray 10 ppm Ethrel as a plant growth regulator at flowering stage. If possible Provide irrigation to cashewnut plant from flowering to fruiting stage. Spraying of 25% cow urine @ 5 liter per plant and drenching of 10 liter of 25% cow urine per cashewnut tree is suggested for increasing yield and size of nuts. If nuts are ready for harvesting, harvest the matured nuts and sun dry for 7 to 8 days to bring down moisture content and then stored in dry places. |

| | | |
|--|-----------------|---|
| Coconut | -- | <ul style="list-style-type: none"> • There is possibility of incidence of rugose spiralling white fly on coconut, Nymphs and adults suck the sap from lower surface of leaves and produce honey dew sugary substance which develop growth of sooty mould fungus. For protection of plant from white fly incidence spray water onto leaves regularly. Also, for control of black fungus spray 1% starch solution on leaves. • Increase in temperature may leads to accelerate evaporation, hence provide irrigation to coconut orchard at 5-6 days interval also use straw mulch to reduce evaporation losses. • Provide shed to the newly planted coconut orchard to protect from sun scorching. |
| Arecanut | - | <ul style="list-style-type: none"> • Increase in temperature may leads to accelerate evaporation, hence provide irrigation to arecanut orchard at 4-5 days interval. |
| Okra | Fruiting | <ul style="list-style-type: none"> • There is possibility of incidence of jassids on okra crop. If incidence is noticed, spray Dimethoate 30%EC@10 ml or Lambda cyhalothrin 5%EC @ 6 ml per 10 liter of water. Also use yellow sticky cards in field to attract sucking pest. |
| <p>This Agro Advisory Bulletin (AAB) is prepared and published with the consultation and recommendation of SMS committees of “Gramin Krishi Mausam Sewa (GKMS)” Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli. For more information contact nearby SAU research station or Agriculture officers of Agriculture Department, Maharashtra state.</p> | | |