

Agromet Advisory Service Bulletin for Raigad District (Issued jointly by GKMS, Dr. B.S. Konkan Krishi Vidyapeeth, & Regional India Meteorological Department, Mumbai) (02358) 282387



No. 20/2020 Date: 09/03/2020 Duration – 5 days

Dr. Prashant Bodake,Head,
Department of Agronomy
9420413255

Dr. Vijay More,Nodal Officer,
Department of Agronomy
9422374001

Dr. Shital Yadav,Technical Officer,
Department of Agronomy
8379901160

Significant past weather for the preceding week (Period -03/03/2020 to 09/03/2020)						eek	Weather Parameters	Weather forecast until 08.30 hrs of 14/03/2020				
03/03	04/03	05/03	06/03	07/03	08/03	09/03		10/03	11/03	12/03	13/03	14/03
-	-	-	-	-	-	-	Rainfall (mm)	0	0	0	0	0
-	-	-	-	-	-	-	Maximum temperature (°C)	29	28	28	29	30
-	-	-	-	-	-	-	Minimum temperature (°C)	22	21	21	21	21
-	-	-	-	-	-	-	Cloud cover (Octa)	0	0	0	0	3
-	-	-	-	-	-	-	Relative Humidity Max. (%)	75	80	76	83	68
-	-	-	-	-	-	-	Relative Humidity Min. (%)	36	47	35	31	33
-	-	-	-	-	-	-	Wind speed (Km/hr)	4	5	4	6	8
-	-	-	-	-	-	-	Wind direction	SSW	WSW	ESE	ENE	ENE
	Rainfall (mm) in last week					Rainfall (mm) from 01/01/2020 to till dated	0 Total Rainfall (mm) in last		st year			
0.0					0.0		5197.2					

Agro-met Advisory
Sky remain clear from 10th to 14th March, 2020.

~	T a.	Sky remain clear from 10 th to 14 th March, 2020.
Crop	Stage	Agro Advise
Summer rice	Panicle	• Maintain optimum water level of 5 cm in rice field where crop is in panicle initiation
	initiation	stage.
Mango	Fruiting	• To minimize the pre-mature fruit drop of mango, apply 150 to 200 liter of water per tree
		after fruit setting (pea size) at 15 days interval for 3 to 4 times also use straw mulch to
		reduce evaporation losses.
		• To improve production and quality of mango fruits, spray 1% Potassium nitrate at pea,
		marble and egg fruit stages.
		• The pre-harvest bagging with newspaper bag of size 25 X 20 cm at marble to egg stage as
		per recommendation of D.B.S.K.K.V. helps to reduce the fruit drop, increases the fruit
		weight, pulp weight, produce spongy tissue free fruit, controls attack of fruit fly on fruits
		and produces spotless fruits of mango.
		• Spraying of 55% cow urine at pea size fruits of mango 3 to 6 sprays at weekly interval as
		per availability is suggested for increasing yield.
Cashewnut	Flowering to	• There is possibility of incidence of tea mosquito bug and thrips on new vegetative flush
	fruiting	of cashewnut, to protect the flush of cashew spray Monocrotophos 36%SL @ 15 ml or
		Lambda cyhalothrin 5% EC @ 6 ml per 10 liter of water.
		• To improve fruit setting and production of cashewnut, spray 10 ppm Ethrel as a plant
		growth regulator at flowering stage. If possible Provide irrigation to cashewnut plant
		from flowering to fruiting stage.
		• Spraying of 25% cow urine @ 5 liter per plant and drenching of 10 liter of 25% cow urine
		per cashewnut tree is suggested for increasing yield and size of nuts.
		• If nuts are ready for harvesting, harvest the matured nuts and sun dry for 7 to 8 days to
		bring down moisture content and then stored in dry places.
Coconut		• There is possibility of incidence of black headed caterpillar in coconut orchard in coastal
		areas. The caterpillar feeds on leaves results into dried up patches on leaf. For control of
		black headed cater pillar, collect and destroy all infected leaves and spray dimethoate
		30%EC @16 ml per 10 liter of water or release larval parasitoids Goniozus nephantidis
		adult @ 3500 nos per hectare.
		• Provide irrigation to coconut orchard at 5-6 days interval also use straw mulch to reduce
		evaporation losses.
Brinjal,	Flowering to	• Provide irrigation to vegetable crops at 6-7 days interval. Also earthing up of soil to give
Tomato, Chilli	fruiting	support to plant.
Water melon	Flowering to	• Provide irrigation to water melon crops at 3-4 days interval regularly as to protect fruits
	fruiting	from cracking.
		• Cover the water melon fruits with paddy straw or grasses to protect fruits from sunlight.
	l	Land Land Land Land Land Land Land Land

This Agro Advisory Bulletin (AAB) is prepared and published with the consultation and recommendation of SMS committees of "Gramin Krishi Mausam Sewa (GKMS)" Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli. For more information contact nearby SAU research station or Agriculture officers of Agriculture Department, Maharashtra

state.	