



Ministry of Earth Science, India Meteorological Department is collaborated with
Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli,
Gramin Krishi Mousam Seva Project,
College of Horticulture, Mulde
Agro Advisory Bulletin For The District Sindhudurg
(Period 10th to 14th March, 2020)



Issue 19/2020

Date 10.03.2020

Duration 5 Days

Actual weather parameters recorded during last week
(Dated 03rd to 09th March, 2020)

Weather Parameters

Forecasted weather parameters for forthcoming 5 day's
(Valid for 10th to 14th March, 2020)

03/03	04/03	05/03	06/03	07/03	08/03	09/03	Date	10/03	11/03	12/03	13/03	14/03	
0.0	0.0	0.0	0.0	0.0	0.0	0.0	Rainfall (mm)	0	0	0	0	0	
36.0	36.0	35.0	33.0	34.0	35.5	35.5	Temp. Maximum (°C)	33	32	32	32	32	
19.0	19.0	18.0	19.0	19.0	18.0	17.5	Temp. Minimum (°C)	19	20	20	19	20	
Cloudy	Clear	Clear	Clear	Cloudy	Clear	Cloudy	AM	Cloud Cover (Octa.)	0	0	1	0	0
Clear	Clear	Clear	Cloudy	Cloudy	Cloudy	Clear	PM						
89	83	88	89	89	83	85	RH - I (%)	66	82	71	63	71	
37	31	44	44	44	37	45	RH - II (%)	38	40	41	33	36	
1.7	1.9	2.6	2.1	1.7	2.2	1.7	Wind Speed (km/hr.)	007	010	009	008	009	
Calm	Calm	Calm	Calm	Calm	Calm	Calm	AM	Wind Direction	148	172	158	86	114
WNW	WNW	WSW	WNW	WNW	WNW	WSW	PM						
Rainfall (mm) in last year (2019)			Rainfall (mm) from 01/01/20 to till dated			Rainfall (mm) from 01/06/20 to till dated			Rainfall (mm) in last week				
4591.0			0.0			0.0			0.0				

General Weather condition in forthcoming 5 days: Wind speed will be increased in forthcoming 5 days also weather remains dry and Clear.

Weather Prediction Based Agro-Advice

Name of Crop	Growth Stage	Advisory
Mango	Flowering to Fruit Set	<p>Due to forecast of increase wind speed in forthcoming week, give support to the newly planted mango seedlings with the help of stick and create shade by using shadenet.</p> <p>Install University recommended methyl eugenol containing 'Rakshak' fruit fly traps @4 traps per ha. to control mango fruit fly.</p> <p>To control mango fruit drop in lateritic rocky soils spraying with 1% Calcium nitrate is recommended.</p> <p>To control mango fruit drop increasing the production and quality of fruits of mango, spraying with 1% Potassium nitrate at pea stage, marble stage and arecanut size stage is recommended.</p> <p>Application of 80-100 litres of water per tree at 7 days interval as per the availability is suggested. Mulching with grass should be done to avoid the water loss in mango orchards in lateritic soil.</p> <p>To decrease fruit drop in mango it is recommended that cover the fruit with paper bags.</p>
Cashewnut	Fruit Set	<p>Due to forecast of increase wind speed in forthcoming week, give support to the newly planted cashewnut seedlings with the help of stick and create shade by using shadenet.</p> <p>To Control of Apple and nut borer of cashew spraying of Diclorovos 76% EC @ 10 ml. per 10 liter of water is advised.</p> <p>To get white colors and whole nut from cashews to process the cashewnut as per University recommendation.</p> <p>Harvest the harvestable and dried cashews from plant and dry it in sunlight.</p> <p>If the incidence of cashew stem rot borer is found to control cashew stem rot borer remove the bark of affected stem with help of 15 mm knife and remove the cashew stem rot borer from stem and soak the stem in to the paste of <i>Cloropyrifos</i> 20% EC @ 10 ml. per 10 liter water is advocated.</p>
Coconut		<p>Due to forecast of increase wind speed in forthcoming week, give support to the newly planted coconut seedlings with the help of stick and create shade by using shadenet.</p> <p>Due to Cloudy & hot weather the incidence of rugose spiraling whitefly is found on coconut. To control establish yellow sticky traps and for removing sooty mould spraying of 1% Starch solution is advised. Also in sever attack spraying of Neem oil 0.5% or 0.005% Imidacloprid is advocated.</p>
Rice	Panicle imitation	<p>Maintain the water level 10 cm for initial 10 days and after 10 days of panicle initiation in rice is advocated.</p> <p>To control blast diseases in rice spraying of <i>Tricyclazole</i> 10 gram or <i>Isoprothionion</i> 10 ml per 10 liter of water is advocated.</p>
Groundnut	flowering to peg formation	<p>Due to forecast of rise in temperature irrigate the groundnut crop with the interval of 12 to 20 days in flowering and pod filling stage.</p> <p>The groundnut crop is in flowering stage and some early varieties is in peg formation stage so drum rolling operation should be carried out to increase number of pods ultimately production will be increased.</p>
Horse gram	Harvesting	In some places Horse gram crop is in harvesting stage. Harvest the mature pods and dry them in sun.

This Agro Advisory Bulletin (AAB) is prepared and published with the consolation and recommendation of SMS committee of "Gramin Krishi Mousam Seva (GKMS)", Regional Fruit Research Station, Vengurle and College of Horticulture, Mulde Dr. B.S.Konkan Krishi Vidyapeeth, Dapoli 416 520 (MS).

Dr. P. C. Haldavanekar
Associate Dean
College of Horticulture, Mulde
Tal. Kudal Dist. Sindhudurg

Dr. P. C. Haldavanekar
Associate Dean and Nodal Officer
College of Horticulture, Mulde
Tal. Kudal Dist. Sindhudurg
(02362-244231,244272)

Dr. Y. C. Muthal
Technical Officer
Gramin Krishi Mousam Seva
College of Horticulture, Mulde
(02362-244231)