

## Agromet Advisory Service Bulletin for Palghar District (Issued jointly by IAAS, Dr. B.S. Konkan Krishi Vidyapeeth, & Regional India Meteorological Department, Mumbai) (02358) 282387



No. 82/2019 Date: 11/10/2019 Duration – 5 days

Dr. Prashant Bodake, Head, Department of Agronomy 9420413255 **Dr. Vijay More,**Nodal Officer,
Department of Agronomy
9422374001

Dr. Shital Yadav, Technical Officer, Department of Agronomy 8379901160

Si	gnificant (Per	-		the prec o 11/10/2	_	eek	Weather Parameters	Weather forecast until 08.30 hrs of 16/10/2019				
05/10	06/10	07/10	08/10	09/10	10/10	11/10		12/10	13/10	14/10	15/10	16/10
0.0	0.0	0.0	0.0	0.0	0.0	0.0	Rainfall (mm)	17	13	5	0	0
32.0	32.4	33.0	33.0	34.0	34.0	34.0	Maximum temperature (°C)	34	34	35	36	35
24.6	25.0	26.0	27.4	27.6	26.0	26.4	Minimum temperature (°C)	24	25	25	25	25
0	0	0	0	0	0	0	Cloud cover (Octa)	8	8	7	5	4
84	84	82	85	86	82	83	Relative Humidity Max. (%)	76	76	76	67	61
72	70	-	62	64	61	-	Relative Humidity Min. (%)	57	59	56	50	49
6.8	1.3	1.6	2.7	2.6	0.7	2.9	Wind speed (Km/hr)	6.7	6.6	6.8	9.5	9.1
NW	S	NNE	NNE	WSW	N	S	Wind direction	ESE	Е	ENE	ENE	ENE
	Rainfa	all (mm)	in last w	eek		F	Rainfall (mm) from 01/01/2019 to till dated	Total Rainfall (mm) in last year			ast year	
0.0							4156.4 2359.6			59.6		

**Agro-met Advisory** 

There is possibility of light to moderate rainfall from 12<sup>th</sup> to 14<sup>th</sup> October, 2019 at isolate places also possibility of increase in maximum temperature and sky will be cloudy.

Extended range rainfall forecast for Konkan division for the period from 11<sup>th</sup> to 17<sup>th</sup> October, 2019 is above normal. According to NDVI, Agriculture vigour is moderate and according to SPI, severely wet condition experienced in Palghar district

		district.
Crop	Stage	Agro Advise
Kharif rice	Grain	• Maintain 5 cm level of water in rice field where late rice varieties are in grain filling stage and
	filling to	drain out water from the field where mid-late rice varieties are in maturity stage.
	maturity	• There is forecast of light to moderate rain, hence by observing rainfall situation harvest matured
	stage	early rice varieties in morning hours only with 'Vaibhav sickle' near to ground level and follow immediate threshing, the threshed grains should be sun dry for 2 to 3 days before storage.
Mango	Vegetative	• In high density (5X5 m or 6X4 m) mango orchard, carryout pruning operation, it includes
		detopping, pruning of cross branches and removing of dead wood. The height of tree in high
		density orchard should be maintained at 80% of row distance.
		• In old mango orchard, carryout centre opening operation which increase penetration of sunlight
		and result in increase in flowering.
		• Due to increase in temperature and humidity there is possibility of incidence of hoppers, midge fly
		and shoot borer on vegetative flush of mango to protect the flush of mango, spray Lambda
		cyhalothrin 5%EC @ 6ml or Quinalphos 25%EC@25 ml per 10 liter of water.
Coconut	Fruiting	• Due to increase in temperature, provide irrigation to coconut orchard.
Sapota	Fruiting	• Follow the weeding operation in sapota orchard, also remove all dried, unproductive and disease
		infected branches, apply bordopaste on cut portion. After cleaning of orchard, spray 1% bordo
		mixture on to tree. For control of seed borer incidence on sapota, after cleaning of orchard, spray
		Profenophos 50% EC @ 15 ml or Indoxacarb 14.5 SC@ 5 ml or Novaluron 10%EC@ 5 ml or
		Deltamethrin 2.8 EC @ 10 ml insecticide per 10 liter of water. Repeat the spray at an interval of a
		month. Harvest all matured fruits before spraying.
	~ .	Due to increase in temperature, provide irrigation to sapota orchard.
Vegetables	Sowing	• For raising of vegetable nursery for rabi season, cultivate the land when it is at field capacity
crop nursery		condition and then prepare raised bed of 3m length x 1 m breadth x 15cm height. Apply 5 kg
		FYM, 35gm urea, 100gm single super phosphate and 25gm muriate of potash per sq. m. For
		protection of seedlings from wilt disease, drench nursery bed with 1% bordomixture before 3-4
		days of sowing. Before sowing, treat the seed with Thiram or Captan fungicide 3 grams per kg of
Milala		seed.
Milch animals/Goat		• Protect the farm animals from increasing temperature.
animais/Goat	11: D	• Provide clean and hygienic water to farm animals also maintain good aeration in farm shed.

This Agro Advisory Bulletin (AAB) is prepared and published with the consultation and recommendation of SMS committees of "Gramin Krishi Mausam Sewa (GKMS)" Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli. For more information contact nearby SAU research station or Agriculture officers of Agriculture Department, Maharashtra state.