



Agromet Advisory Service Bulletin for Raigad District
(Issued jointly by GKMS, Dr. B.S. Konkan Krishi Vidyapeeth,
& Regional India Meteorological Department, Mumbai)
(02358) 282387



No. 21/2020

Date: 13/03/2020

Duration – 5 days

Dr. Prashant Bodake,
Head,
Department of Agronomy
9420413255

Dr. Vijay More,
Nodal Officer,
Department of Agronomy
9422374001

Dr. Shital Yadav,
Technical Officer,
Department of Agronomy
8379901160

Significant past weather for the preceding week (Period –07/03/2020 to 13/03/2020)							Weather Parameters	Weather forecast until 08.30 hrs of 18/03/2020				
07/03	08/03	09/03	10/03	11/03	12/03	13/03		14/03	15/03	16/03	17/03	18/03
-	-	-	-	-	-	-	Rainfall (mm)	0	0	0	0	0
-	-	-	-	-	-	-	Maximum temperature (°C)	29	30	31	32	32
-	-	-	-	-	-	-	Minimum temperature (°C)	19	21	22	22	21
-	-	-	-	-	-	-	Cloud cover (Octa)	0	0	2	0	0
-	-	-	-	-	-	-	Relative Humidity Max. (%)	57	39	38	58	49
-	-	-	-	-	-	-	Relative Humidity Min. (%)	27	17	16	21	22
-	-	-	-	-	-	-	Wind speed (Km/hr)	8	5	4	3	5
-	-	-	-	-	-	-	Wind direction	ESE	ENE	ESE	E	E
Rainfall (mm) in last week						Rainfall (mm) from 01/01/2020 to till dated			Total Rainfall (mm) in last year			
0.0						0.0			5197.2			

Agro-met Advisory

Sky remain clear from 14th to 18th March, 2020.

Crop	Stage	Agro Advise
Summer rice	Panicle initiation	<ul style="list-style-type: none"> Maintain optimum water level of 5 cm in rice field where crop is in panicle initiation stage. There is possibility of incidence of stem borer in rice, if incidence is noticed, apply Quinalphos 5% granules @ 15 kg/ha or Carbofuron 3 % granules @ 16.5 kg/ha when there is sufficient moisture in the field.
Lablab bean, Cowpea	Pod development stage	<ul style="list-style-type: none"> There is possibility of incidence of sucking pest on lanlab bean and cowpea etc. pulse crops., If incidence is noticed, spray Dimethoate 30%EC@12 ml or Lambda cyhalothrin 5% EC @ 6 ml per 10 liter of water.
Mango	Fruiting	<ul style="list-style-type: none"> To minimize the pre-mature fruit drop of mango, apply 150 to 200 liter of water per tree after fruit setting (pea size) at 15 days interval for 3 to 4 times also use straw mulch to reduce evaporation losses. To improve production and quality of mango fruits, spray 1% Potassium nitrate at pea, marble and egg fruit stages. The pre-harvest bagging with newspaper bag of size 25 X 20 cm at marble to egg stage as per recommendation of D.B.S.K.K.V. helps to reduce the fruit drop, increases the fruit weight, pulp weight, produce spongy tissue free fruit, controls attack of fruit fly on fruits and produces spotless fruits of mango. Spraying of 55% cow urine at pea size fruits of mango 3 to 6 sprays at weekly interval as per availability is suggested for increasing yield.
Cashewnut	Flowering to fruiting	<ul style="list-style-type: none"> There is possibility of incidence of tea mosquito bug and thrips on new vegetative flush of cashewnut, to protect the flush of cashew spray Monocrotophos 36%SL @ 15 ml or Lambda cyhalothrin 5% EC @ 6 ml per 10 liter of water. There is possibility of incidence of tea mosquito bugs and thrips on the inflorescence of cashewnut, to protect the cashew inflorescence, spray Profenophos 50% EC @ 10 ml per 10 liter of water and for control of incidence of tea mosquito buds and thrips during fruit bearing stage of cashewnut, spray Lambda cyhalothrin 5% EC @ 6 ml per 10 liter of water. (insecticide is not under label claim) If nuts are ready for harvesting, harvest the matured nuts and sun dry for 7 to 8 days to bring down moisture content and then stored in dry places.
Leafy vegetables	Vegetative	<ul style="list-style-type: none"> There is possibility of incidence of leaf spot disease in leafy vegetables. For control of disease, spray 0.1% carbenazim @ 1 gm/liter or 0.25% copper oxychloride @2.5 gm/liter of water at 15 days interval.
Water melon	Flowering to fruiting	<ul style="list-style-type: none"> Provide irrigation to water melon crops at 3-4 days interval regularly as to protect fruits from cracking. Cover the water melon fruits with paddy straw or grasses to protect fruits from sunlight. Install cue lure 'Rakshak' trap @ 4 nos. per hectare at the time of initiation of in watermelon crop for effective control of fruit fly. Collect and destroy all infected fruits.
<p>This Agro Advisory Bulletin (AAB) is prepared and published with the consultation and recommendation of SMS committees of "Gramin Krishi Mausam Sewa (GKMS)" Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli. For more information contact nearby SAU research station or Agriculture officers of Agriculture Department, Maharashtra state.</p>		