

Agromet Advisory Service Bulletin for Raigad District (Issued jointly by IAAS, Dr. B.S. Konkan Krishi Vidyapeeth, & Regional India Meteorological Department, Mumbai) (02358) 282387



No. 74/2019 Date: 13/09/2019 Duration – 5 days

Dr. Prashant Bodake,Head,
Department of Agronomy
9420413255

Dr. Vijay More,Nodal Officer,
Department of Agronomy
9422374001

Dr. Shital Yadav,Technical Officer,
Department of Agronomy
8379901160

Sig	gnificant (Per	past wea iod –07/0				eek	Weather Parameters	Wea	Weather forecast until 08.30 hrs of 18/09/2019			
07/09	08/09	09/09	10/09	11/09	12/09	13/09		14/09	15/09	16/09	17/09	18/09
85.0	84.2	12.3	67.2	28.5	21.2	106.4	Rainfall (mm)	52	25	23	23	2
28.4	27.6	28.8	-	-	-	-	Maximum temperature (°C)	29	30	31	29	29
24.6	23.5	24.5	-	-	-	-	Minimum temperature (°C)	24	24	25	24	26
6.0	8.0	8.0	-	-	-	-	Cloud cover (Octa)	8	7	5	6	7
88	95	87	-	-	-	-	Relative Humidity Max.(%)	95	94	94	94	89
92	82	89	-	-	-	-	Relative Humidity Min. (%)	86	84	75	82	81
5.9	6.7	5.6	-	-	-	-	Wind speed (Km/hr)	9	5	5	4	4
ESE	SSE	N	-	-	-	-	Wind direction	251	201	208	220	220
	Rainf	all (mm)	in last w	eek		F	Rainfall (mm) from 01/01/2019 to till dated		Total Rainfall (mm) in last year			
	404.8						4268.6	3558.3				

Agro-met Advisory

There is possibility of moderate to heavy rainfall from 14th to 17th September, 2019. Sky will be cloudy.

Extended r	ange rainfall f	forecast for Konkan division for the period of 13 th to 19 th September, 2019 is above normal.
Crop	Stage	Agro Advise
Kharif rice	Panicle initiation to flowering	 Maintain the optimum water level of 5-10 cm in the rice crop which is in panicle initiation to flowering stage. Apply 3rd split dose of nitrogen @ 43 kg urea/ha when intensity of rainfall is low to the midlate varieties rice crop which is in flowering stage.
Mango	Vegetative	 Due to forecast of increase in temperature which is congenial for vegetative flush in mango, for protection of vegetative flush, spray Lambda cyhalothrin 5%EC@ 6ml or Quinalphos 25% EC@25 ml + Bavistin 20 g per 10 liter of water.
Coconut		 Apply 750 gms urea and 667 gms of muriate of potash as 2nd split dose for 5 year old and above coconut palm. There is possibility of incidence of bud rot disease due to heavy rain,if incidence noticed spray coconut plant with 1% bordomixture. Clean old coconut orchard by removing all dried leaves and other dead waste to maintain good sanitation in orchard. There is possibility of bud rot due to heavy rain, if incidence noticed spray 1% bordomixture on to the crown region.
Vegetable crops	Fruiting	 Due to forecast of cloudy weather and increase in humidity, there is possibility of incidence of downy mildew in cucurbitaceous vegetables, if incidence noticed, remove and destroy infected leaves and take a spray of Mancozeb or Zineb 2.5 gm /lit of water at an interval of 10 to 15 days. There is possibility of incidence of red pumpkin beetle on cucumber, bitter gourd, snake gourd, bottle gourd, red pumpkin and ridge gourd etc. cucurbitaceous crops. If incidence is noticed, spray Dimethoate 30%EC@ 15 ml or Malathion 50%EC @ 20 ml per 10 liters of water.
Fruit crop nursery		• To prevent the incidence of fungal disease in fruit crop nursey, drenching of 50 ml of fungicidal solution (0.2% Carbendazim @ 2 gm/lit of water) per plant is advocated.
Milch animals/Goat /Poultry		 For young goat, provide 200 to 250 gms Concentrate + 3 to 4 kg green fodder + 1.5 kg roughages per day. For boiler poultry bird 1st three week provide boiler starter and 4 to 6 week old bird provide boiler finisher as per their daily requirement.

Note: spraying should be taken in dry period during day.

This Agro Advisory Bulletin (AAB) is prepared and published with the consultation and recommendation of SMS committees of "Gramin Krishi Mausam Sewa (GKMS)" Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli. For more information contact nearby SAU research station or Agriculture officers of Agriculture Department, Maharashtra state.