

Agromet Advisory Service Bulletin for Raigad District (Issued jointly by GKMS, Dr. B.S. Konkan Krishi Vidyapeeth, & Regional India Meteorological Department, Mumbai) (02358) 282387



No. 100/2019 Date: 13/12/2019 Duration – 5 days

Dr. Prashant Bodake, Head, Department of Agronomy 9420413255 **Dr. Vijay More,**Nodal Officer,
Department of Agronomy
9422374001

Dr. Shital Yadav,Technical Officer,
Department of Agronomy
8379901160

Significant past weather for the preceding week (Period -07/12/2019 to 13/12/2019)							Weather Parameters	Wea	Weather forecast until 08.30 hrs of 18/12/2019			
07/12	08/12	09/12	10/12	11/12	12/12	13/12		14/12	15/12	16/12	17/12	18/12
-	-	-	-	-	-	-	Rainfall (mm)	0	0	0	0	0
-	-	-	-	-	-	-	Maximum temperature (°C)	30	30	29	29	29
-	-	-	-	-	-	-	Minimum temperature (°C)	21	21	20	20	20
-	-	-	-	-	-	-	Cloud cover (Octa)	5	3	4	2	2
-	-	-	-	-	-	-	Relative Humidity Max.(%)	75	66	67	68	71
-	-	-	-	-	-	-	Relative Humidity Min. (%)	43	40	43	42	40
-	-	-	-	-	-	-	Wind speed (Km/hr)	7	6	5	7	8
-	-	-	-	-	-	-	Wind direction	NE	ENE	ENE	ENE	Е
	Rainfall (mm) in last week						Rainfall (mm) from 01/01/2019 to till dated	Total Rainfall (mm) in last ye		st year		
0.0							5222.1		3558.3			

Agro-met Advisory

Sky will be partly cloudy from 14th to 16th December 2019.

Extended range rainfall forecast for Konkan division for the period from 13th to 19th December, 2019 is below normal. According to NDVI, Agriculture vigour is good and according to SPI, mildly wet condition experienced in Raigad district

		re vigour is good and according to SPI, mildly wet condition experienced in Raigad district.
Crop	Stage	Agro Advise
Lablab bean	Vegetative	• If incidence of aphids is noticed on lablab bean, spray Dimethoate 30%EC@12 ml per 10 liter of water.
Horse gram	Vegetative	• If incidence of leaf folder is noticed on horse gram, spray Dimethoate 30%EC@20 ml or Profenophos 50%EC @ 20 ml per 10 liter of water.
White Onion	Nursery	 If incidence of thrips is noticed on onion, spray Quinalphos 25 ml per 10 liters of water. For control of weeds in onion crop nursery, do weeding operation at 8-10 days interval also provide irrigation.
Mango	Vegetative	• There is possibility of incidence of hoppers and anthracnose disease on vegetative flush of mango. For protection of vegetative flush from hoppers and diseases, spray deltamethrin 2.8%EC @ 10 ml + fungicide combination (carbendazim 12% + mancozeb 63%) @ 10 gm per 10 liter in water.
Cashewnut	Flowering	• There is possibility of incidence of tea mosquito bugs on the inflorescence of cashewnut, to protect the cashew inflorescence, spray Profenophos 50% EC @10 ml per 10 liter of water.
Coconut	-	• There is possibility of incidence of black headed caterpillar in coconut orchard in coastal areas. The caterpillar feeds on leaves results into dried up patches on leaf. For control of black headed cater pillar, collect and destroy all infected leaves and spray dimethoate 30%EC @16 ml per 10 liter of water or release larval parasitoids Goniozus nephantidis adult @ 3500 nos per hectare.
Arecanut	-	• For 3-year-old arecanut palm apply 2 nd split dose 160 g urea and 125 g muriate of potash per tree. by digging circular ring about 1 meter from base of tree and fill the circular ring with soil after application of fertilizer. Apply 1/3 rd and 2/3 rd of abovementioned fertilizers dose per tree for 1 and 2-years old arecanut plantation, respectively.
Chilli	Vegetative	• There is possibility of incidence of hoppers, white files and aphids in chilli crop. If incidence is noticed, spray dimethoate 30%EC @10 ml per 10 liter of water. Also use yellow sticky cards in field to attract the sucking pests.

This Agro Advisory Bulletin (AAB) is prepared and published with the consultation and recommendation of SMS committees of "Gramin Krishi Mausam Sewa (GKMS)" Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli. For more information contact nearby SAU research station or Agriculture officers of Agriculture Department, Maharashtra state.