



**Agromet Advisory Service Bulletin for Palghar District**  
(Issued jointly by IAAS, Dr. B.S. Konkan Krishi Vidyapeeth,  
& Regional India Meteorological Department, Mumbai)  
(02358) 282387



No. 83/2019

Date: 15/10/2019

Duration – 5 days

**Dr. Prashant Bodake,**  
Head,  
Department of Agronomy  
9420413255

**Dr. Vijay More,**  
Nodal Officer,  
Department of Agronomy  
9422374001

**Dr. Shital Yadav,**  
Technical Officer,  
Department of Agronomy  
8379901160

Significant past weather for the preceding week (Period –09/10/2019 to 15/10/2019)							Weather Parameters	Weather forecast until 08.30 hrs of 20/10/2019				
09/10	10/10	11/10	12/10	13/10	14/10	15/10		16/10	17/10	18/10	19/10	20/10
0.0	0.0	0.0	0.0	0.0	0.0	0.0	Rainfall (mm)	0	0	0	2	8
34.0	34.0	34.0	33.0	34.0	35.0	-	Maximum temperature (°C)	35	34	35	30	32
27.6	26.0	26.4	25.0	25.4	25.6	-	Minimum temperature (°C)	25	25	26	24	23
0	0	0	0	0	0	-	Cloud cover (Octa)	3	8	4	8	8
86	82	83	87	84	85	-	Relative Humidity Max. (%)	61	64	65	73	79
64	61	60	51	56	51	-	Relative Humidity Min. (%)	48	52	53	64	62
2.6	0.7	2.9	1.9	2.1	1.3	-	Wind speed (Km/hr)	10.7	10.9	11.2	11.6	12.5
WSW	N	S	NNE	NNW	NNE	-	Wind direction	ENE	ENE	E	E	E
Rainfall (mm) in last week						Rainfall (mm) from 01/01/2019 to till dated		Total Rainfall (mm) in last year				
0.0						4156.4		2359.6				

**Agro-met Advisory**

There is possibility of light rainfall on 19<sup>th</sup> and 20<sup>th</sup> October, 2019 at isolate places also possibility of increase in maximum temperature and sky will be cloudy.

Extended range rainfall forecast for Konkan division for the period from 18<sup>th</sup> to 24<sup>th</sup> October, 2019 is above normal. According to NDVI, Agriculture vigour is moderate and according to SPI, severely wet condition experienced in Palghar district.

Crop	Stage	Agro Advise
<b>Kharif rice</b>	<b>Grain filling to maturity stage</b>	<ul style="list-style-type: none"> <li>• Maintain 5 cm level of water in rice field where late rice varieties are in grain filling stage and drain out water from the field where mid-late rice varieties are in maturity stage.</li> <li>• There is forecast of light rain, hence by observing rainfall situation harvest matured early rice varieties in morning hours only with 'Vaibhav sickle' near to ground level and follow immediate threshing, the threshed grains should be sun dry for 2 to 3 days before storage.</li> <li>• There is possibility of incidence of army worm in early rice varieties, hence by observing rainfall situation harvest matured early rice varieties.</li> </ul>
<b>Mango</b>	<b>Vegetative</b>	<ul style="list-style-type: none"> <li>• Due to forecast for increase in temperature, provide irrigation to young mango graft either by conventional or with drip irrigation method also apply straw or black polythene mulch around tree.</li> <li>• In high density (5X5 m or 6X4 m) mango orchard, carryout pruning operation, it includes detopping, pruning of cross branches and removing of dead wood. The height of tree in high density orchard should be maintained at 80% of row distance.</li> <li>• In old mango orchard, carryout centre opening operation which increase penetration of sunlight and result in increase in flowering.</li> <li>• Due to increase in temperature and humidity, there is possibility of incidence of hoppers, midge fly and shoot borer on vegetative flush of mango to protect the flush of mango, spray Lambda cyhalothrin 5%EC @ 6ml or Quinalphos 25%EC@25 ml per 10 liter of water.</li> </ul>
<b>Coconut</b>	<b>Fruiting</b>	<ul style="list-style-type: none"> <li>• Due to increase in temperature, provide irrigation to coconut orchard.</li> </ul>
<b>Sapota</b>	<b>Fruiting</b>	<ul style="list-style-type: none"> <li>• Follow inter cultivation operation in sapota orchard. collect and destroy all infected fruits and dried leaves.</li> </ul>
<b>Vegetables crop nursery</b>	<b>Sowing</b>	<ul style="list-style-type: none"> <li>• For raising of vegetable nursery for rabi season, cultivate the land when it is at field capacity condition and then prepare raised bed of 3m length x 1 m breadth x 15cm height. Apply 5 kg FYM, 35gm urea, 100gm single super phosphate and 25gm muriate of potash per sq. m. For protection of seedlings from wilt disease, drench nursery bed with 1% bordomixture before 3-4 days of sowing. Before sowing, treat the seed with Thiram or Captan fungicide 3 grams per kg of seed.</li> <li>• Provide irrigation to fruit crop and vegetable nursery regularly.</li> </ul>
<b>Poultry</b>	-	<ul style="list-style-type: none"> <li>• Protect poultry birds from increasing temperature.</li> </ul>

**This Agro Advisory Bulletin (AAB) is prepared and published with the consultation and recommendation of SMS committees of "Gramin Krishi Mausam Sewa (GKMS)" Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli.**  
**For more information contact nearby SAU research station or Agriculture officers of Agriculture Department, Maharashtra state.**