



**Agromet Advisory Service Bulletin for Thane District**  
(Issued jointly by GKMS, Dr. B.S. Konkani Krishi Vidyapeeth,  
& Regional India Meteorological Department, Mumbai)  
(02358) 282387



No. 101/2019

Date: 17/12/2019

Duration – 5 days

**Dr. Prashant Bodake,**  
Head,  
Department of Agronomy  
9420413255

**Dr. Vijay More,**  
Nodal Officer,  
Department of Agronomy  
9422374001

**Dr. Shital Yadav,**  
Technical Officer,  
Department of Agronomy  
8379901160

Significant past weather for the preceding week (Period –11/12/2019 to 17/12/2019)							Weather Parameters	Weather forecast until 08.30 hrs of 22/12/2019				
11/12	12/12	13/12	14/12	15/12	16/12	17/12		18/12	19/12	20/12	21/12	22/12
0	0	0	0	0	0	-	Rainfall (mm)	0	0	0	0	0
35.0	32.9	32.5	34.1	35.4	34.4	-	Maximum temperature (°C)	32	32	33	33	31
22.7	21.2	20.3	19.4	19.7	20.7	-	Minimum temperature (°C)	21	21	22	22	22
-	-	-	-	-	-	-	Cloud cover (Octa)	2	3	2	6	7
100	97	99	100	86	98	-	Relative Humidity Max. (%)	67	65	62	60	59
77	73	78	73	79	78	-	Relative Humidity Min. (%)	43	38	37	35	40
2	1	3	3	3	2	-	Wind speed (Km/hr)	8	7	6	4	4
NE	SE	NE	ENE	NE	NE	-	Wind direction	NE	NE	NE	ENE	E

**Agro-met Advisory**

Sky will be partly cloudy from 18<sup>th</sup> to 22<sup>nd</sup> December 2019.

Extended range rainfall forecast for Konkani division for the period from 13<sup>th</sup> to 19<sup>th</sup> December, 2019 is below normal.  
According to NDVI, Agriculture vigour is good and according to SPI, mildly wet condition experienced in Thane district.

Crop	Stage	Agro Advise
Lablab bean	Vegetative	• If incidence of aphids is noticed on lablab bean, spray Dimethoate 30%EC@12 ml per 10 liter of water.
Mango	Vegetative	• There is possibility of incidence of hoppers and anthracnose disease on vegetative flush of mango. For protection of vegetative flush from hoppers and diseases, spray deltamethrin 2.8%EC @ 10 ml + fungicide combination (carbendazim 12% + mancozeb 63%) @ 10 gm per 10 liter in water.
Cashewnut	Flowering	• There is possibility of incidence of tea mosquito bugs on the inflorescence of cashewnut, to protect the cashew inflorescence, spray Profenophos 50% EC @10 ml per 10 liter of water.
Coconut	-	• There is possibility of incidence of black headed caterpillar in coconut orchard in coastal areas. The caterpillar feeds on leaves results into dried up patches on leaf. For control of black headed caterpillar, collect and destroy all infected leaves and spray dimethoate 30%EC @16 ml per 10 liter of water or release larval parasitoids <i>Goniozus nephantidis</i> adult @ 3500 nos per hectare.
Banana	Vegetative	• If incidence of bunchy top disease is noticed on banana, remove and destroy infected plant. Since the aphid act as vector for bunchy top disease, if incidence of aphid is notice in banana orchard spray 0.05 % dimethoate insecticide at 15 days interval.
Tomato	Vegetative	• Provide support to tomato crop. • Apply second split dose of 43kg urea per acre to tomato crop at one month after transplanting.
Brinjal	Vegetative	• Apply second split dose of 43kg urea per acre to brinjal crop at one month after transplanting.
Marigold	Vegetative	• For control of weeds in marigold flower crop, do weeding operation after 15-20 days of transplanting also earthing up of soil to give support to plant.
Aster	Vegetative	• For control of weeds in aster flower crop, do weeding operation after 15-20 days of transplanting also earthing up of soil to give support to plant.
Milch Animals/ Goat /Poultry	--	• Provide clean and hygienic drinking water to farm animals. • Protect the animals /poultry birds from low temperature by providing curtains and electric bulbs as per the need.

**This Agro Advisory Bulletin (AAB) is prepared and published with the consultation and recommendation of SMS committees of “Gramin Krishi Mausam Sewa (GKMS)” Dr. Balasaheb Sawant Konkani Krishi Vidyapeeth, Dapoli.**  
For more information contact nearby SAU research station or Agriculture officers of Agriculture Department, Maharashtra state.