



**Agromet Advisory Service Bulletin for Thane District**  
(Issued jointly by GKMS, Dr. B.S. Konkan Krishi Vidyapeeth,  
& Regional India Meteorological Department, Mumbai)  
(02358) 282387



No. 06/2020

Date: 21/01/2020

Duration – 5 days

**Dr. Prashant Bodake,**  
Head,  
Department of Agronomy  
9420413255

**Dr. Vijay More,**  
Nodal Officer,  
Department of Agronomy  
9422374001

**Dr. Shital Yadav,**  
Technical Officer,  
Department of Agronomy  
8379901160

Significant past weather for the preceding week (Period –15/01/2020 to 21/01/2020)							Weather Parameters	Weather forecast until 08.30 hrs of 26/01/2020				
15/01	16/01	17/01	18/01	19/01	20/01	21/01		22/01	23/01	24/01	25/01	26/01
0	0	0	0	0	0	-	Rainfall (mm)	0	0	0	0	0
29.5	26.5	28.5	33.1	34.8	35.5	-	Maximum temperature (°C)	33	34	34	34	33
17.8	16.4	13.7	15.5	18	19.6	-	Minimum temperature (°C)	17	18	18	18	17
-	-	-	-	-	-	-	Cloud cover (Octa)	0	0	1	6	5
86	97	80	96	88	95	-	Relative Humidity Max. (%)	56	50	56	56	47
67	61	69	54	57	65	-	Relative Humidity Min. (%)	35	32	31	30	23
2	3	3	4	3	3	-	Wind speed (Km/hr)	6	7	7	6	7
NE	NE	NE	NE	NE	NE	-	Wind direction	ENE	ENE	ENE	ENE	NE

**Agro-met Advisory**  
**Sky will remain mainly cloudy from 25<sup>th</sup> to 26<sup>th</sup> January, 2020.**

Crop	Stage	Agro Advise
<b>Groundnut</b>	<b>Vegetative</b>	<ul style="list-style-type: none"> <li>There is possibility of incidence of aphids on Groundnut crop which develop growth of black fungus on leaves and inflorescence. If incidence is noticed, spray Dimethoate 30%EC@12 ml per 10 liter of water.</li> </ul>
<b>Lablab bean</b>	<b>Flowering</b>	<ul style="list-style-type: none"> <li>There is possibility of incidence of pod borer on lablab bean crop which initially feed on buds and then on tender pods. If incidence is noticed, collect and destroy all infected pods and spray Quinalphos 25% EC @ 20ml of Dimethoate 30% EC@12 ml per 10 liter of water. Install birds' perches into field.</li> <li>Provide irrigation to lablab bean crop where crop is in flowering stage.</li> </ul>
<b>Mango</b>	<b>flower bud initiation to flowering</b>	<ul style="list-style-type: none"> <li>To protect the flower bud of mango from hoppers, thrips and powdery mildew diseases, spray Lambda cyhalothrin 5%EC @ 6 ml + hexaconazole @ 5 ml per 10 liter in water at the time of flower bud initiation as a second spray of mango blossom protection schedule.</li> <li>As per blossom protection schedule for mango crop, take a third spray of Imidacloprid 17.8% SL @ 6 ml per 10 liter of water before the flower opening (15 days after 2<sup>nd</sup> spray) to avoid the adverse effect on pollinators. Also add Hexaconazole 5% @ 5 ml or wettable Sulphur 80% @ 20 gm per 10 liter in water for control of powdery mildew disease.</li> </ul>
<b>Sapota</b>	<b>Fruiting</b>	<ul style="list-style-type: none"> <li>Apply second split dose of 5 kg FYM, 150 g urea, 450 g single super phosphate and 150 g muriate of potash per tree to year old sapota plant by band placement around the tree just inside the spread. Apply fertilizer dose every year by multiplying year with first year dose upto first 20 years and after 20 years, apply 100 kg FYM, 3 kg urea, 9 kg single super phosphate and 3 kg muriate of potash per tree thereafter.</li> </ul>

**This Agro Advisory Bulletin (AAB) is prepared and published with the consultation and recommendation of SMS committees of "Gramin Krishi Mausam Sewa (GKMS)" Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli. For more information contact nearby SAU research station or Agriculture officers of Agriculture Department, Maharashtra state.**