



Agromet Advisory Service Bulletin for Ratnagiri District
(Issued jointly by GKMS, Dr. B.S. Konkan Krishi Vidyapeeth,
& Regional India Meteorological Department, Mumbai)
(02358) 282387



No. 07/2020

Date: 24/01/2020

Duration – 5 days

Dr. Prashant Bodake,
Head,
Department of Agronomy
9420413255

Dr. Vijay More,
Nodal Officer,
Department of Agronomy
9422374001

Dr. Shital Yadav,
Technical Officer,
Department of Agronomy
8379901160

Significant past weather for the preceding week (Period –18/01/2020 to 24/01/2020)							Weather Parameters	Weather forecast until 08.30 hrs of 29/01/2020				
18/01	19/01	20/01	21/01	22/01	23/01	24/01		25/01	26/01	27/01	28/01	29/01
0.0	0.0	0.0	0.0	0.0	0.0	0.0	Rainfall (mm)	0	0	0	0	0
26.0	30.0	32.2	33.4	32.0	33.6	34.0	Maximum temperature (°C)	34	33	31	30	29
11.0	13.5	15.0	15.5	16.5	15.0	15.0	Minimum temperature (°C)	20	21	19	18	17
0	0	0	0	0	0	0	Cloud cover (Octa)	1	5	3	5	1
93	95	93	94	92	93	93	Relative Humidity Max. (%)	61	57	64	85	86
60	50	51	59	52	42	-	Relative Humidity Min. (%)	28	27	30	39	48
3.7	2.6	2.0	2.5	2.9	2.7	3.3	Wind speed (Km/hr)	5	5	3	4	5
Calm	Calm	Calm	Calm	Calm	Calm	Calm	Wind direction	ENE	E	ESE	SE	E
Rainfall (mm) in last week							Rainfall (mm) from 01/01/2020 to till dated	Total Rainfall (mm) in last year				
0.0							0.0	5130.9				

Agro-met Advisory

There is possibility of decrease in maximum and minimum temperature and sky will remain partly cloudy from 26th to 28th January, 2020.

Crop	Stage	Agro Advise
Lablab bean	Flowering	<ul style="list-style-type: none"> There is possibility of incidence of pod borer on lablab bean crop which initially feed on buds and then on tender pods. If incidence is noticed, collect and destroy all infected pods and spray Quinalphos 25% EC @ 20ml of Dimethoate 30% EC@12 ml per 10 liter of water. Install birds' perches into field. Provide irrigation to lablab bean crop where crop is in flowering to pod filling stage.
Mango	flower bud initiation to flowering	<ul style="list-style-type: none"> To protect the flower bud of mango from hoppers, thrips and powdery mildew diseases, spray Lambda cyhalothrin 5%EC @ 6 ml + hexaconazole @ 5 ml per 10 liter in water at the time of flower bud initiation as a second spray of mango blossom protection schedule. As per blossom protection schedule for mango crop, take a third spray of Imidacloprid 17.8% SL @ 6 ml per 10 liter of water before the flower opening (15 days after 2nd spray) to avoid the adverse effect on pollinators. Also add Hexaconazole 5% @ 5 ml or wettable Sulphur 80% @ 20 gm per 10 liter in water for control of powdery mildew disease.
Cashewnut	Flowering to fruiting	<ul style="list-style-type: none"> There is possibility of incidence of tea mosquito bugs and thrips on the inflorescence of cashewnut, to protect the cashew inflorescence, spray Profenophos 50% EC @10 ml per 10 liter of water and for control of incidence of tea mosquito buds and thrips during fruit bearing stage of cashewnut, spray Lambda cyhalothrin 5% EC @6 ml per 10 liter of water. (insecticide is not under label claim)
Coconut	--	<ul style="list-style-type: none"> Provide irrigation to first four years old coconut orchard at 6 to 7 days interval and for above four years old orchard provide irrigation at 5 to 10 days interval. To control the attack of red palm weevil on coconut, collect and destroy the grubs from whole appear on infected trunk. Apply bordopaste to infected part. Install pheromone trap 2 nos. per hectare in orchard.

This Agro Advisory Bulletin (AAB) is prepared and published with the consultation and recommendation of SMS committees of "Gramin Krishi Mausam Sewa (GKMS)" Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli. For more information contact nearby SAU research station or Agriculture officers of Agriculture Department, Maharashtra state.