



Ministry of Earth Science, India Meteorological Department is collaborated with
 Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli,
Gramin Krishi Mousam Seva Project,
 College of Horticulture, Mulde
 Agro Advisory Bulletin For The District Sindhudurg
 (Period 28th December, 2019 to 1st January, 2020)



Issue 104/2019

Date 27.12.2019

Duration 5 Days

Actual weather parameters recorded during last week (Dated 21 st to 27 th December, 2019)							Weather Parameters		Forecasted weather parameters for forthcoming 5 day's (Valid for 28 th December, 2019 to 1 st January, 2020)				
21/12	22/12	23/12	24/12	25/12	26/12	27/12	Date	28/12	29/12	30/12	31/12	01/01	
0.0	0.0	0.0	0.0	0.0	0.0	0.0	Rainfall (mm)	0	0	2	4	0	
35.0	36.0	36.0	35.0	35.5	35.5	34.0	Temp. Maximum (°C)	32	32	32	32	30	
19.0	20.0	20.0	20.0	21.0	21.5	22.0	Temp. Minimum (°C)	23	23	23	21	20	
Clear	Clear	Clear	Cloudy	Cloudy	Cloudy	Clear	AM	Cloud Cover (Octa.)	4	3	1	0	0
Clear	Clear	Cloudy	Clear	Cloudy	Cloudy	Clear	PM						
91	91	91	84	84	90	91	RH - I (%)	80	82	83	90	85	
37	37	42	41	42	47	47	RH - II (%)	48	45	47	50	52	
0.9	1.6	1.2	0.7	1.3	1.1	0.6	Wind Speed (km/hr.)	007	006	005	005	005	
Calm	Calm	Calm	Calm	Calm	Calm	Calm	AM	Wind Direction	123	81	63	71	88
WNW	WSW	WSW	ENE	ENE	WSW	ENE	PM						

Rainfall (mm) in last year (2018)	Rainfall (mm) from 01/01/19 to till dated	Rainfall (mm) from 01/06/19 to till dated	Rainfall (mm) in last week
3269.2	4591.0	4591.0	0.0

General Weather condition in forthcoming 5 days: Light rain is predicted in forthcoming 5 days in the district and Atmosphere remain partly cloudy.

Weather Prediction Based Agro-Advice

Name of Crop	Growth Stage	Advisory
Wayangani Rice	Rabbi sowing	In Rabbi rice, maintain the water level 2-3 cm for first 8-10 days and increase the level up to 5 cm.
Horse gram and Cowpea	Seedling to Branching	The incidence of aphids and leaf miner may be noticed on pulse crops. For their control spraying with 10 ml Dimethoate per 10 litres of water at an interval of 15-20 days is suggested.
Groundnut Sunflower and Mustard	Rabbi sowing	Light irrigation to groundnut crop three days after sowing. Irrigation interval should be 10-12 days for this stage. Also irrigate the sunflower and mustard crop timely. Gap filling should be done in groundnut crop.
Mango	Flushing	Due to extended post monsoon during this year, mango trees is not acrossed with water stress and the flowering stage of mango is delayed also due to frequent cyclones in the last month atmosphere remain cloudy and minimum temperature is not decreased and temperature and atmosphere is favorable for mango flowering. Protect the new flushing in mango may be observed simultaneously the incidence of anthracnose and powdery mildew diseases may found on mango. In cloudy weather the host is expected to emerge is large quantity. Shade and high humidity are favorable for their multiplication. Due to heavy leaf flush on mango and such condition usually can make a huge attack on mango hopper on inflorescence leaf and fruits. To control spraying of Thiophenate methyl and Carbendanzim 1% @ 1 gram per liter water is advocated. Incidence of Mango Hoper, Thrips and Scale insect may be come on New flush of mango if found the incidence to control mango hopper application of insecticide like Deltamethrin 2.8% EC (0.9 per liter of water) or Thiomethoxam 25% WDG (0.1 gram per liter of water) is advised.
Cashewnut	Flowering	Current Climatic conditions are favorable for cashew borer. If such incidence is notified on inflorescence of cashew then spraying of 50 per cent Profenophos 10 ml/10 litres of water and if fruiting is their then spraying of 5 per cent lamdasayhalothrin 6ml/10 liters of water is advocated. If the incidence of cashew stem rot borer is found to control cashew stem rot borer remove the bark of affected stem with help of 15 mm knife and remove the cashew stem rot borer from stem and soak the stem in to the paste of <i>Cloropyrifhos</i> 20% EC @ 10 ml. per 10 liter water is advocated. To Control of Apple and nut borer of cashew spraying of Diclorovos 76% EC @ 10 ml. per 10 liter of water is advised.
Vegetable		Installation of Cue-lure containing Rakshak fruit fly traps @ 4 traps/ha for control of fruit fly in gourd crops is advocated.
Dairy Animals, Goat, Sheep Poultry		Protect the animals from cold during the night time as the night temperatures are decreasing. For protection of poultry birds from cold and cold winds during night, provide curtain/clothing from outside of poultry shed. Provide proper heat to small chicks of poultry with the help of electric bulbs.

This Agro Advisory Bulletin (AAB) is prepared and published with the consolation and recommendation of SMS committee of "Gramin Krishi Mausam Sewa (GKMS)", Regional Fruit Research Station, Vengurle and College of Horticulture, Mulde Dr. B.S.Konkan Krishi Vidyapeeth, Dapoli 416 520 (MS).

Dr. P.C. Haldavanekar
 Associate Dean
 College of Horticulture, Mulde
 Tal. Kudal Dist. Sindhudurg

Dr. P.C. Haldavanekar
 Associate Dean and Nodal Officer
 College of Horticulture, Mulde
 Tal. Kudal Dist. Sindhudurg
 (02362-244231,244272)

Dr. Y. C. Muthal
 Technical Officer
 Gramin Krishi Mousam Seva
 College of Horticulture, Mulde
 (02362-244231)