



Agromet Advisory Service Bulletin for Raigad District
(Issued jointly by GKMS, Dr. B.S. Konkan Krishi Vidyapeeth,
& Regional India Meteorological Department, Mumbai)
(02358) 282387



No. 09/2020

Date: 31/01/2020

Duration – 5 days

Dr. Prashant Bodake,
Head,
Department of Agronomy
9420413255

Dr. Vijay More,
Nodal Officer,
Department of Agronomy
9422374001

Dr. Shital Yadav,
Technical Officer,
Department of Agronomy
8379901160

Significant past weather for the preceding week (Period –25/01/2020 to 31/01/2020)							Weather Parameters	Weather forecast until 08.30 hrs of 05/02/2020				
25/01	26/01	27/01	28/01	29/01	30/01	31/01		01/02	02/02	03/02	04/02	05/02
-	-	-	-				Rainfall (mm)	0	0	0	0	0
-	-	-	-				Maximum temperature (°C)	27	28	29	27	28
-	-	-	-				Minimum temperature (°C)	17	18	19	17	18
-	-	-	-				Cloud cover (Octa)	0	1	3	4	5
-	-	-	-				Relative Humidity Max. (%)	59	54	74	71	62
-	-	-	-				Relative Humidity Min. (%)	31	26	27	28	29
-	-	-	-				Wind speed (Km/hr)	6	6	5	3	8
-	-	-	-				Wind direction	NE	E	E	ENE	NE
Rainfall (mm) in last week							Rainfall (mm) from 01/01/2020 to till dated	Total Rainfall (mm) in last year				
0.0							0.0	5197.2				

Agro-met Advisory

During next five days the maximum temperature will be in the range of 27 to 29°C and minimum temperature in the range of 17 to 19 °C and sky remain partly cloudy from 3rd to 5th February, 2020.

Crop	Stage	Agro Advise
Rice	Seedling	• Maintain optimum water level of 2.5- 5 cm upto 30 days after transplanting of rice seedling.
Lablab bean	Pod stage	• There is possibility of incidence of pod borer on lablab bean crop which initially feed on buds and then on tender pods. If incidence is noticed, collect and destroy all infected pods and spray Quinalphos 25% EC @ 20ml of Dimethoate 30% EC@12 ml per 10 liter of water. Install birds' perches into field. • Provide irrigation to lablab bean crop where crop is in flowering stage.
Chickpea	Flowering	• There is possibility of incidence of pod borer on chickpea crop, for protection install pheromone trap and birds's perches @ 5 -10 nos. per hectare.
Mango	Flowering	• To protect the flower bud of mango from hopppers, thrips and powdery mildew diseases, as per blossom protection schedule for mango crop, take a third spray of Imidacloprid 17.8% SL @ 6 ml per 10 liter of water before the flower opening (15 days after 2 nd spray) to avoid the adverse effect on pollinators. Also add Hexaconazole 5% @ 5 ml or wettable Sulphur 80% @ 20 gm per 10 liter in water for control of powdery mildew disease.
Cashewnut	Fruiting	• There is possibility of incidence of tea mosquito bugs and thrips on the fruits of cashewnut, to protect the cashew during fruit bearing stage, spray Lambda cyhalothrin 5% EC @6 ml per 10 liter of water. (insecticide is not under label claim)
Brinjal	Fruiting	• There is possibility of incidence of shoot and fruit borer on Brinjal, if incidence is noticed, collect and destroy all infected shoots and fruits and spray Fenvalerate @ 10 ml or Deltamethrin 208%EC @ 10 ml per 10 liter of water.
Water melon	Fruiting	• There is possibility of incidence of red pumpkin beetle on water melon, if incidence is noticed, spray Dimethoate 30%EC @15 ml or Malathion 50%EC @ 20 ml per 10 liter of water.
Milch Animals	--	• Protect the farm animals from low temperature by providing curtains and electric bulbs as per the need.
Poultry	--	• Protect the poultry birds from low temperature by providing curtains and electric bulbs as per the need.

This Agro Advisory Bulletin (AAB) is prepared and published with the consultation and recommendation of SMS committees of "Gramin Krishi Mausam Sewa (GKMS)" Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli. For more information contact nearby SAU research station or Agriculture officers of Agriculture Department, Maharashtra state.