



Agromet Advisory Service Bulletin for Thane District
(Issued jointly by GKMS, Dr. B.S. Konkan Krishi Vidyapeeth,
& Regional India Meteorological Department, Mumbai)
(02358) 282387



No. 79/2020

Date: 01/10/2020

Duration – 5 days

Dr. Prashant Bodake,
Head,
Department of Agronomy
9420413255

Dr. Vijay More,
Nodal Officer,
Department of Agronomy
9422374001

Dr. Shital Yadav,
Technical Officer,
Department of Agronomy
8379901160

Significant past weather for the preceding week (Period –25/09/2020 to 01/10/2020)							Weather Parameters	Weather forecast until 08.30 hrs of 06/10/2020				
25/09	26/09	27/09	28/09	29/09	30/09	01/10		02/10	03/10	04/10	05/10	06/10
3.0	4.0	0.0	18.0	0.0	-	-	Rainfall (mm)	13	18	20	18	9
32.0	32.6	34.2	33.8	33.0	-	-	Maximum temperature (°C)	33	33	33	33	34
24.2	-	24.0	24.0	25.0	-	-	Minimum temperature (°C)	25	25	25	25	26
4	4	3	5	4	-	-	Cloud cover (Octa)	5	7	4	3	3
94	-	89	98	90	-	-	Relative Humidity Max. (%)	90	90	91	93	93
80	82	74	83	78	-	-	Relative Humidity Min. (%)	69	68	68	72	71
0.0	0.0	0.0	0.0	0.0	-	-	Wind speed (Km/hr)	4	4	8	5	5
Calm	Calm	Calm	Calm	Calm	-	-	Wind direction	SSW	SW	W	SSW	S

Weather summary

Weather forecast	As per the forecast received from Regional Meteorological Centre Mumbai, there is possibility of moderate rainfall from 2 nd to 5 th October, 2020 and light rainfall 6 th October, 2020 at a few places over Thane district. Also, sky will remain partly cloudy.
Warning	Thunderstorm accompanied with lighting likely to occur on 3 rd to 4 th October, 2020 at isolated places in Thane district.
ERFS	As per ERFS products normal rainfall may occur in Konkan division during 9 th to 15 th October, 2020.

Crop	Stage	Agromet Advisory based on weather forecast
Kharif rice	Flowering to grain filling (late varieties), grain filling (midlate varieties) and maturity (early varieties)	<ul style="list-style-type: none"> Due to forecast of light to moderate rainfall from 3rd to 6th October, 2020, harvest the matured early rice varieties during morning hours, stored threshed grains in protective shed after immediate threshing. Dry threshed grains in protective shed. Maintain the optimum water level of 5-10 cm in late and water of 5 cm in midlate rice varieties field. Apply 3rd split dose of nitrogen @ 28 kg urea per acre to the late varieties of rice which is in flowering. Due to forecast of intermittent rainfall and increase in temperature, there is possibility of incidence of army worm in early rice varieties, hence immediate harvest the matured early rice varieties. Due to forecast of humid weather condition during morning hours, there is possibility of incidence of rice earhead bug where late and midlate rice varieties are in milking stage. If incidence of rice earhead bug is observed above threshold level (1 adult/hill), spray Lambda cyhalothrin 5%EC @ 5 ml or Deltamethrin 2.8%EC @ 9 ml per 10 liter of water. (Note: insecticide is not under label claim).
Mango	Vegetative	<ul style="list-style-type: none"> Due to forecast of very light rainfall and increase in temperature which favours congenial condition for vegetative flush in mango, for protection of vegetative flush of mango from hoppers and midge fly incidence, spray Deltamethrin 2.8%EC @ 9 ml per 10 liter of water. Before insecticide application, clean orchard by removing weeds, diseased infected branches.
Coconut	Fruiting	<ul style="list-style-type: none"> Due to forecast for increase in temperature, provide shed to newly planted coconut orchard (upto 2 years old) as to protect leaf from sunburn.
Sapota	Fruiting	<ul style="list-style-type: none"> Follow inter cultivation operation in sapota orchard. collect and destroy all infected fruits and dried leaves.
Vegetable crops	Sowing	<ul style="list-style-type: none"> For raising brinjal, chilli, tomato and other vegetable crop nursery for rabi season, cultivate the land at field capacity condition and mix well decomposed FYM @500 gms per guntha in to soil at the time of bed preparation. Prepare raised bed of 3 m length x 1 m breadth x 15 cm height. At the time of sowing apply 35 gm urea, 100 gm single super phosphate and 25 gm muriate of potash per sq. m. Before sowing, treat the seed with thiram fungicide @ 3 gm/kg of seed or Trichoderma @ 5 gm/kg for protection from seed borne diseases. for seed treatment of Trichoderma use water to prepare paste to adhere like seed coat.

Note : Spray insecticide or fungicide solution and apply fertilizers when intensity of rainfall is low. Mix sticker into spraying solution.

This Agro Advisory Bulletin (AAB) is prepared and published with the consultation and recommendation of SMS committees of "Gramin Krishi Mausam Sewa (GKMS)" Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli. For more information contact nearby SAU research station or Agriculture officers of Agriculture Department, Maharashtra

state.