



Agromet Advisory Service Bulletin for Raigad District
(Issued jointly by GKMS, Dr. B.S. Konkan Krishi Vidyapeeth,
& Regional India Meteorological Department, Mumbai)
(02358) 282387



No. 53/2020

Date: 03/07/2020

Duration – 5 days

Dr. Prashant Bodake,
Head,
Department of Agronomy
9420413255

Dr. Vijay More,
Nodal Officer,
Department of Agronomy
9422374001

Significant past weather for the preceding week (Period –27/06/2020 to 03/07/2020)							Weather Parameters	Weather forecast until 08.30 hrs of 08/07/2020				
27/06	28/06	29/06	30/06	01/07	02/07	03/07		04/07	05/07	06/07	07/07	08/07
0.0	0.0	7.0	0.0	6.2	8.0	27.2	Rainfall (mm)	249	98	79	89	93
33.2	34.0	33.8	33.2	-	-	-	Maximum temperature (°C)	29	27	28	27	27
26.4	26.2	26.0	25.4	-	-	-	Minimum temperature (°C)	24	24	24	24	24
4	6	4	4	-	-	-	Cloud cover (Octa)	7	8	7	7	7
91	91	79	89	-	-	-	Relative Humidity Max. (%)	97	96	95	95	96
66	67	71	78	-	-	-	Relative Humidity Min. (%)	80	94	91	90	93
3.2	4.3	2.8	5.3	-	-	-	Wind speed (Km/hr)	15	11	16	14	10
ENE	NE	NNE	Calm	-	-	-	Wind direction	SW	SW	SW	SW	SSW
Rainfall (mm) in last week				Rainfall (mm) from 01/01/2020 to till dated				Total Rainfall (mm) in last year				
48.4				455.4				5197.2				

Agro-met Advisory

There is possibility of rainfall at most places over Raigad district from 4th to 8th July, 2020

Warning: Heavy to very heavy rainfall likely to occur at few place and extremely heavy rainfall at isolated places on 4th July 2020 and heavy to very heavy rainfall from 5th to 8th July, 2020 at isolated places.

As per ERFS products above normal rainfall may occur in Konkan division during 08th to 14th July 2020.

Crop	Stage	Agro Advise
Kharif rice	Transplanting	<ul style="list-style-type: none"> Due to forecast of heavy rain, make arrangement for drain out excess water from rice field and maintain optimum water level of 2.5 to 5cm upto 30 days after transplanting. Carry out transplanting of the rice seedlings with 5 to 6 leaves with height of 12 to 15 cm and 20-21 days old seedling in case of early rice varieties and 27 to 30 days old seedling in case of midlate and late rice varieties. Prepare field for transplanting by puddling. As there is forecast for heavy to very heavy rainfall on 4th and 5th July, postponed the basal application of fertilizers to rice at the time of puddling. Apply when intensity of rainfall is low. Transplant 2-3 rice seedlings per hill in case of straight varieties and one seedling per hill in case of hybrid at a spacing of 15 x 15 cm (early) or 20 x 15 cm (midlate and late varieties). follow upright and shallow (2.5 to 3.5 cm) transplanting of seedling. In SIRA method of rice cultivation, incorporate 2 tonnes per acre glyricida leaves at the time of puddling. Transplant improved or hybrid rice seedlings at a controlled spacing (15-25 cms). place urea-DAP briquette at depth of 7.5 to 10 cms in between four hill.
Finger millet	Transplanting	<ul style="list-style-type: none"> Due to forecast of heavy rain, make arrangement for drain out excess water from finger millet nursery. It is advise to undertake transplanting of finger millet seedlings when intensity of rainfall is low. Seedlings of 30 days old are consider ideal for transplanting. Transplant 2 finger millet seedlings at spacing of 20x15 cm by 'Thomba' method. Prepare Thomba across the slope. Apply 35 kg urea and 100 kg single super phosphate per acre in thomba at the time of transplanting.
Coconut, Arecanut	Fruiting	<ul style="list-style-type: none"> Due to forecast of heavy rain, make arrangement for drain out excess water from coconut and arecanut orchard
Vegetable crops	Vegetative	<ul style="list-style-type: none"> Due to forecast of heavy rain, make arrangement for drain out excess water from vegetable crop field.
Turmeric	Vegetative	<ul style="list-style-type: none"> Make arrangement for drainage of excess water from turmeric field.
Milch animals	-	<ul style="list-style-type: none"> Provide clean and hygienic drinking water to farm animals also maintain good aeration in farm shed. Protect the farm animals from rain water.

Note : Spray insecticide or fungicide solution and apply fertilizers when intensity of rainfall is low. Mix sticker into spraying solution.

This Agro Advisory Bulletin (AAB) is prepared and published with the consultation and recommendation of SMS committees of "Gramin Krishi Mausam Sewa (GKMS)" Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli. For more information contact nearby SAU research station or Agriculture officers of Agriculture Department, Maharashtra state.