


Ministry of Earth Science, India Meteorological Department is collaborated with Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli,

Gramin Krishi Mousam Sewa,

College of Horticulture, Mulde





Date 06.10.2020 Issue 79/2020 **Duration 5 Days** Actual weather parameters recorded during last week Forecasted weather parameters for forthcoming 5 day's **Weather Parameters** (Dated 29th September to 05th October, 2020) (Valid for 07th to 11th October, 2020) 29/09 30/09 01/10 02/10 03/10 04/10 05/10 07/10 08/10 09/10 10/10 11/10 Date 12.0 7.2 18.3 6.5 8.4 5 0.0 10.6 9 6 0 13 Rainfall (mm) 32.5 33 34 34 34 33 32.0 32.5 33.0 33.0 33.0 33.0 Temp. Maximum (°C) 23.0 23.0 23.0 23.0 22.0 23.0 22.5 24 25 24 24 24 Temp. Minimum (°C) Cloudy Cloudy Clear Clear Clear Cloudy Cloudy AM **Cloud Cover** 7 8 2 4 8 Cloudy Cloudy Cloudy Cloudy Cloudy PM Cloudy Cloudy (Octa.) 92 94 94 93 90 96 96 98 98 96 95 93 RH - I (%) 71 57 **75** 84 **72** 60 60 67 67 78 65 56 RH - II (%) 0.1 0.2 0.2 0.2 0.2 0.2 0.2 004 005 005 003 003 Wind Speed (km/hr.) Calm Calm Calm Calm Calm Calm Calm **AM** Wind Easterly/ Easterly/ East East South West South East South East Calm Calm Calm Calm **WSW** Calm **WSW** PM **Direction** Rainfall (mm) from 01/01/20 to till dated Rainfall (mm) from 01/06/20 to till dated Rainfall (mm) in last year (2019) Rainfall (mm) in last week 4591.0 4551.2

General Weather conditions & warnings in forthcoming 5 days:

October 11, 2020.

In Sindhudurg district during the next five days from 07 to 11 October, 2020 there is a possibility of light rain in isolated places between 07 October to 10 October and light rain in scattered places on 11 October. At the same time, the weather is likely to remain humid and mainly cloudy with a slight rise in temperature. This forecast has been made by the Regional Meteorological Center, Mumbai.

Warning: In Sindhudurg district, light rain with gusts of wind, thunderstorms accompanied with lightning is expected in scattered places on

For control of anthracnose on new flush of mango, application of Propineb @ 20 gm. or Thiophenate methyl @ 10 gm. or

Weather Forecast Based Agro-Advice				
Name of Crop	Growth Stage	Advisory		
Paddy		There was light to moderate rain last week and light rain is expected in the next five days.		
		Considering the possibility of very light rains in the next 5 days, the harvest of early paddy varieties should be started.		
		If early paddy varieties are ready to be harvested, arrangements should be made to remove water from the field 8 to 10 days before		
	Early	harvest. No pesticide / fungicide spray or dusting should be done on paddy 8 days before harvesting.		
	varieties:-	In some places early rice varieties are ready for harvest. The paddy should be cut near the ground from morning to noon at the		
	Harvesting	place where the paddy is ready for harvesting.		
		The harvested paddy crop should be followed by threshing and keeping the grains in a dry and safe place for drying.		
	Mid late & late	Vaibhav sickle developed by the university should be used for paddy harvesting so that the paddy is cut near the ground.		
	varieties :-	Arrangements should be made to keep the paddy crop straw in a safe place after threshing. So that the straw will not get wet in the		
	Flowering to	rain and good fodder will be available to the animals.		
	grain filling	In the district late varieties are in flowering to grain filling stage and in such places 5 to 10 cm water level should be kept.		
		In some places in the district where late rice varieties are in the stage of flowering to grain filling, apply third dose of nitrogen		
		fertilizer (435 gm urea per guntha area) at crop flowering stage. Fertilizer application should be done in the absence of rainy		
		conditions.		
Finger Millet	Panicle	According to the weather forecast, light rains is expected from October 7 to 11, 2020. In the morning, the earheads of the early		
	initiation	Finger Millet crop should be cut with a sickle and spread in a dry and safe place for drying and threshed after drying.		
Groundnut		The groundnut crop is in flowering stage and some early varieties is in peg formation stage so drum rolling operation should be		
		carried out to increase number of pods ultimately production will be increased.		
	Flowering to	At some places groundnut crop is in peg initiation stage, so proper care of the crop should be taken. If the incidence of aphids is		
	Peg formation	observed in groundnut crop, then farmers are suggested to apply Dimethoate @ 10 ml per 10 litres of water for its control.		
		If the incidence of Tikka disease is noticed on groundnut crop, then farmers are advocated to apply Carbendazim @ 10 gms or		
		Dithane M-45 @ 20 gms per 10 litres of water for its control.		
		Keep the paclobutrazol applied trees free from weeds by weeding or applying weedicide Gramoxone. Sanitation of Mango orchard		
Mango		is advocated.		
	New flush	Due to the sudden and intermittent rain, rise in temperature and cloudy weather the incidence of pest may occur on the new flush		
		on mango so protect the new flush from mango hoppers, leaf eating caterpillars, shoot borer and midge fly. If incidence is noticed,		
		spraying with Dimethoate 25 EC @ 12 ml or Monocrotophos 36 WSC @ 11 ml per 10 litres of water as per need is advocated.		

Carbendazim @ 10 gm. per 10 liter of water as per need is advocated.			per need is advocated.		
		For control of dieback disease of mango, spraying with copper oxychloride 0.3 per cent (30 gms/10 litres of water) or 1% Bordeaux			
		mixture is advocated. If there is oozing of gum from matured branches, then scrap the infected branch and apply paste of Copper			
		oxychloride or Bordeaux mixture to the affected portion.			
		Cutting diseased, dried branches, Loranthus par	rasite and apply 1 per cent Bordeaux mixture to the cut surface is advocated.		
		At time of spraying sticky substance like sticker	should be mix with pesticide/fungicide solution @ 1 ml/liter of water.		
Cashew nut	-	In a large cashew orchard, clean the orchard by removing weeds and shrubs.			
		Due to the sudden and intermittent rain, rise in temperature and cloudy weather Cashew orchards are getting new leaves flush in			
		the district Cashew growers should be vigilant to prevent the infestation of Tea mosquito bug, In case of infestation, monocrotophos			
		36% EC @ 15 ml or lambda cyhalothrin 5% @ 6 ml. Spray in 10 liters of water.			
		If the incidence CSRB is noticed in cashew, remove the grubs with the help of 15 mm chisel and soak the stem in to the paste of			
		fipronil 5% SC @ 2 ml per liter of water (5 liter of water is sufficient for one tree) is advocated.			
		To control Dieback disease on cashews, cut the branches and apply 1% Bordeaux paste on it. After the end of rainfall season,			
		spray 1% Bordeaux mixture all over the garden.			
Coconst	Fruit	Newly planted coconut seedlings should be shaded with shade net for the first 2 years as the temperature may rise.			
Coconut	Development				
Arecanut	Fruit	Newly planted arecanut seedlings should be shaded with shade net for the first 2 years as the temperature may rise.			
	Development	Koleroga (fruit drop) disease of Arecanut, spraying of 1% Bordo mixture in the orchard is suggested.			
Vegetable	Fruiting	Protect the vegetables crops from pests like leaf eating caterpillars, aphids etc. by spraying 2 ml of Malathion or 1.5 ml of			
Vegetable		Dimethoate per litre of water for their control.			
Doin: Animal	-	As light rain is expected in scattered places with thunderstorms accompanied with lightning on October 11, farmers should be			
Dairy Animal, Goat		careful while working in the fields to avoid lightning. Also, to protect the dairy animals released from grazing from lightning, the dairy			
		animals should be kept in a safer place.			
This Agro Advisory Bulletin (AAB) is prepared and published with the consolation and recommendation of SMS committee of "Gramin Krishi Mausam Sewa (GKMS)", Regional Fruit Research					
Station, Vengurle and College of Horticulture, Mulde Dr. B.S.Konkan Krishi Vidyapeeth, Dapoli 416 520 (MS).					
		P.C. Haldavanekar Dean and Nodal Officer	Dr. Y. C. Muthal		
	Agro-Meteor	rological Field Unit (AMFU),	Technical Officer		
	Gramin	Krishi Mousam Sewa,	Agro-Meteorological Field Unit (AMFU), Gramin Krishi Mousam Sewa.		
		of Horticulture, Mulde	College of Horticulture, Mulde		
		udal Dist. Sindhudurg	(02362-244231)		
(02362-244231,244272)					