



# AGROMET ADVISORY SERVICE BULLETIN FOR THANE DISTRICT

(Issued jointly by GKMS, Dr. B.S. Konkani Krishi Vidyapeeth,  
& India Meteorological Department)



Ph.No. : (02358) 282387

Email : dpl.amfu@gmail.com

No. 03/2021

Date: 08/01/2021

Duration – 5 days

**Dr. Prashant Bodake,**  
Head,  
Department of Agronomy  
9420413255

**Dr. Vijay More,**  
Nodal Officer,  
Department of Agronomy  
9422374001

**Dr. Shital Yadav,**  
Technical Officer,  
Department of Agronomy  
8379901160

Significant past weather for the preceding week (Period –02/01/2021 to 08/01/2021)							Weather Parameters	Weather forecast until 08.30 hrs of 13/01/2021				
(Source: Agromet observatory, T.B.I.A., Thane)								(Source: Regional Meteorological Centre, Mumbai)				
02/01	03/01	04/01	05/01	06/01	07/01	08/01		09/01	10/01	11/01	12/01	13/01
0.0	0.0	0.0	0.0	0.0	-	-	Rainfall (mm)	12	9	0	0	0
-	31.8	-	33.0	33.2	-	-	Max.Temp. (°C)	31	32	32	33	34
21.2	19.0	18.0	21.0	19.2	-	-	Min.Temp. (°C)	20	21	21	22	23
0	0	0	4	0	-	-	Cloud cover (Octa)	4	7	4	5	7
46	74	62	77	83	-	-	Max. RH (%)	77	74	71	62	57
-	70	-	66	71	-	-	Min. RH (%)	51	49	51	37	37
0.0	0.0	0.0	0.0	0.0	-	-	Wind speed(Km/hr)	7	6	6	7	6
Calm	Calm	Calm	Calm	Calm	-	-	Wind direction	E	NE	ENE	ENE	ENE

### Weather summary/alert

<b>Weather forecast</b>	As per the forecast received from Regional Meteorological Centre Mumbai, there is possibility of light rainfall from 9 <sup>th</sup> to 10 <sup>th</sup> January, 2021 at isolated places and sky remain cloudy from 9 <sup>th</sup> to 13 <sup>th</sup> January, 2021 over Thane district.
<b>Warning</b>	Thunderstorm accompanied with lightning is likely to occur from 8 <sup>th</sup> to 9 <sup>th</sup> January, 2021 at isolated places in Thane district.

### Agromet advisory based on weather forecast

Crop	Stage	Agro advisory
<b>Mango</b>	Vegetative flush	<ul style="list-style-type: none"> <li>• Due to forecast for light rainfall from 8<sup>th</sup> to 10<sup>th</sup> January, 2021 and possibility for cloudy weather condition till 13<sup>th</sup> January 2021, there is possibility for anthracnose disease on mango vegetative flush and inflorescence, for protection against anthracnose disease, spray carbendazim 12% + mancozeb 63% combination fungicide @ 10 gm per 10 liter in water after 10<sup>th</sup> January, 2021.</li> <li>• Due to forecast for light rain with cloudy weather condition, there is possibility incidence of hoppers and powdery mildew disease on flower bud stage in mango. To protect the flower bud from pest and powdery mildew diseases, spray Lambda cyhalothrin 5%EC @ 6 ml + hexaconazole 5% @ 5 ml or wettable Sulphur 80% @ 20 gm per 10 liter in water during clear weather.</li> <li>• The recommended dose of insecticides is applicable for manually operating sprayer.</li> </ul>
<b>Cashewnut</b>	Flowering to fruiting	<ul style="list-style-type: none"> <li>• Due to forecast for light rainfall from 8<sup>th</sup> to 10<sup>th</sup> January, 2021 and possibility for cloudy weather condition till 13<sup>th</sup> January 2021, there is possibility for incidence of anthracnose fungal disease on vegetative flush of cashewnut, to control the disease spray, 1% bordomixture or Carbendazim 12% + Mancozeb 63% combination fungicide @ 10 gms per 10 liter of water after 10<sup>th</sup> January, 2021.</li> <li>• Due to forecast for light rain with cloudy weather condition, there is possibility of incidence of tea mosquito bugs and thrips on the inflorescence of cashewnut, to protect the cashew inflorescence, spray Profenophos 50% EC @10 ml per 10 liter of water during clear weather.</li> <li>• There is possibility of incidence of tea mosquito bugs and thrips on the</li> </ul>

		fruits of cashewnut, to protect the cashew during fruit bearing stage, spray Lambda cyhalothrin 5% EC @6 ml or Acetamiprid 20%SP @ 5 gms per 10 liter of water during clear weather. (insecticide is not under label claim).
<b>Chickpea</b>	Flowering	<ul style="list-style-type: none"> <li>• Due to forecast for light rainfall from 8<sup>th</sup> to 10<sup>th</sup> January, 2021 and cloudy weather condition till 13<sup>th</sup> January 2021, there is possibility for incidence of pod borer in chickpea. For management of pest install pheromone trap @2 trap per acre and install bird's perches. If incidence is noticed spray 0.5% neem seed kernel extract or azadiracthin 300 PPM @ 5 ml per liter of water.</li> </ul>
<b>Lablab bean/ Cowpea</b>	Vegetative to flowering	<ul style="list-style-type: none"> <li>• Due to forecast for light rainfall from 8<sup>th</sup> to 10<sup>th</sup> January, 2021 and cloudy weather condition till 13<sup>th</sup> January 2021, there is possibility of incidence of aphids,hoppers on lablab, cowpea crop which develop into growth of black fungus on leaves and inflorescence. If incidence is noticed, spray Dimethoate 30%EC@12 ml per 10 liter of water during clear weather.</li> </ul>
<b>Chilli, Brinjal, Tomato</b>	Flowering to fruiting	<ul style="list-style-type: none"> <li>• Due to forecast for light rainfall from 8<sup>th</sup> to 10<sup>th</sup> January, 2021 and possibility for cloudy weather condition till 13<sup>th</sup> January 2021, there is possibility of incidence of anthracnose disease in chilli, brinjal, tomato vegetable crops, if incidence is noticed, spray Mancozeb 20 gm per 10 liter of water.</li> </ul>
<b>Cabbage</b>	Vegetative	<ul style="list-style-type: none"> <li>• Due to forecast for light rainfall from 8<sup>th</sup> to 10<sup>th</sup> January, 2021 and cloudy weather condition till 13<sup>th</sup> January 2021, there is possibility of incidence of anthracnose disease in cabbage crop. If incidence is noticed, spray Copper oxychloride 25 gm of per 10 liters of water during clear weather.</li> </ul>
<b>Cucurbitaceous crop</b>	Flowering to fruiting	<ul style="list-style-type: none"> <li>• Due to forecast for light rainfall from 8<sup>th</sup> to 10<sup>th</sup> January, 2021 and possibility for cloudy weather condition till 13<sup>th</sup> January 2021, there is possibility of incidence of downy mildew and powdery mildew diseases in cucumber, snake gourd, bottle gourd, red pumpkin, ridge gourd, bitter gourd, water melon cucurbitaceous vegetables, if incidence of downy mildew is noticed, remove and destroy infected leaves and spray Mancozeb 75% WP @ 25 gm or Coper oxychloride 50%WP @ 20 gms per 10 liters of water during clear weather at an interval of 10 to 15 days and for control of powdery mildew disease spray Carbendazim 10 gm or Hexaconazole 5%EC @ 5 ml per 10 liter of water during clear weather at an interval of 10 to 15 days.</li> </ul>
<p>This Agro Advisory Bulletin (AAB) is prepared and published with the consultation and recommendation of SMS committees of "Gramin Krishi Mausam Sewa (GKMS)" Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli. For more information contact nearby SAU research station or Agriculture officers of Agriculture Department, Maharashtra state.</p>		