



**Agromet Advisory Service Bulletin for Raigad District**  
(Issued jointly by GKMS, Dr. B.S. Konkan Krishi Vidyapeeth,  
& Regional India Meteorological Department, Mumbai)  
(02358) 282387



No. 90/2020

Date: 10/11/2020

Duration – 5 days

**Dr. Prashant Bodake,**  
Head,  
Department of Agronomy  
9420413255

**Dr. Vijay More,**  
Nodal Officer,  
Department of Agronomy  
9422374001

**Dr. Shital Yadav,**  
Technical Officer,  
Department of Agronomy  
8379901160

Significant past weather for the preceding week (Period –04/11/2020 to 10/11/2020)							Weather Parameters	Weather forecast until 08.30 hrs of 15/11/2020				
04/11	05/11	06/11	07/11	08/11	09/11	10/11		11/11	12/11	13/11	14/11	15/11
0.0	-	-	-	-	-	-	Rainfall (mm)	0	0	0	0	0
34.8	-	-	-	-	-	-	Maximum temperature (°C)	34	33	35	36	37
18.4	-	-	-	-	-	-	Minimum temperature (°C)	20	20	23	24	24
4	-	-	-	-	-	-	Cloud cover (Octa)	0	0	4	2	1
94	-	-	-	-	-	-	Relative Humidity Max. (%)	56	54	67	75	71
42	-	-	-	-	-	-	Relative Humidity Min. (%)	31	30	37	48	46
0.9	-	-	-	-	-	-	Wind speed (Km/hr)	9	9	8	7	6
Calm	-	-	-	-	-	-	Wind direction	E	ENE	E	E	ENE
Rainfall (mm) in last week				Rainfall (mm) from 01/01/2020 to till dated				Total Rainfall (mm) in last year				
0.0				3089.4				5197.2				

**Weather summary**

<b>Weather forecast</b>	<b>As per the forecast received from Regional Meteorological Centre Mumbai, there is possibility of clear weather with increase in maximum and minimum temperature from 11<sup>th</sup> to 15<sup>th</sup> November, 2020 over Raigad district.</b>
-------------------------	---

Crop	Stage	Agromet Advisory based on weather forecast
<b>General advice</b>	-	<ul style="list-style-type: none"> <li>Due to increase in rate of evaporation, provide irrigation to irrigated fruit crops, nurseries and vegetable crops.</li> </ul>
<b>Kharif rice</b>	<b>Maturity</b>	<ul style="list-style-type: none"> <li>Sun dry the threshed grain for 2 to 3 days before storage.</li> </ul>
<b>Finger millet</b>	<b>Maturity</b>	<ul style="list-style-type: none"> <li>Sun dry the threshed grain for 2 to 3 days before storage.</li> </ul>
<b>Cowpea</b>	<b>Land preparation and Sowing</b>	<ul style="list-style-type: none"> <li>Plough the field for cultivation of cowpea and incorporate FYM or compost @ 2 tonnes/acre at the time of last ploughing. prepare check basin of 4 X 3 m size leaving furrow of 30 cm for irrigation between alternate two beds. Sow seed at a spacing of 30 X 15 cm. Apply 65 gms urea, 450 gms single Super Phosphate and 80 gms of muriate of potash per check basin before sowing.</li> <li>Konkan sadabahar and Konkan safed are the varieties recommended by D.B.S.K.K.V., for cultivation in Konkan region. Use 6-8 kg seed /acre.</li> <li>Treat the seed with Thiram fungicide @3 gm/kg or Trichoderma 5 gms/kg and with Rhizobium and PSB biofertilizers @ 25 gms per kg of seed before sowing.</li> </ul>
<b>Chickpea</b>	<b>Land preparation and Sowing</b>	<ul style="list-style-type: none"> <li>For cultivation of Chickpea, plough the soil after harvest of kharif rice when it is in field capacity. Incorporate 2 tonne/acre FYM or compost at the time of ploughing. Sow seed at a spacing of 30 x 10 cm. apply 540 gms urea and 3 kg Single Super Phosphate per guntha at the time of sowing. Provide light irrigation after sowing.</li> <li>Treat the seed with Thiram fungicide @3 gm/kg or Trichoderma 5 gms/kg and with Rhizobium and PSB biofertilizers @ 25 gms per kg of seed before sowing.</li> </ul>
<b>Mango</b>	<b>Vegetative</b>	<ul style="list-style-type: none"> <li>For protection of vegetative flush of mango from hoppers and midge fly incidence, spray Deltamethrin 2.8%EC @ 9 ml per 10 liter of water.</li> <li>Situation of clear sky and increase in temperature may leads to accelerate evaporation, hence provide irrigation to newly planted mango orchard @ 30 liters of water at 7 days interval (1 years old), 15 days interval (2 years old) and one-month interval to 3 years old tree. New growth below graft union should be removed regularly.</li> </ul>
<b>Cashewnut</b>	<b>Flowering</b>	<ul style="list-style-type: none"> <li>There is congenial weather situation of flower bud opening in cashewnut, to protect the cashew inflorescence from tea mosquito bug and thrips incidence, spray Profenophos 50% EC @10 ml per 10 liter of water. (The insecticide is not under label claim).</li> <li>To improve fruit setting and production of cashewnut, take 2<sup>nd</sup> spray of 10 PPM ethrel at the flowering stage.</li> <li>Situation of clear sky and increase in temperature may leads to accelerate evaporation, hence provide irrigation to newly planted cashewnut orchard @ 15 liters of water at 15 days intervals. New growth below graft union should be removed regularly.</li> </ul>
<b>Coconut</b>	<b>Fruiting</b>	<ul style="list-style-type: none"> <li>Due to clear sky and increase in temperature accelerate evaporation, provide irrigation to coconut orchard at 8-10 days interval.</li> </ul>

		<ul style="list-style-type: none"> <li>To control the attack of red palm weevil on coconut, collect and destroy the grubs from whole appear on infected trunk. Apply bordopaste to infected portion. Install pheromone trap 2 nos. per hectare in orchard.</li> </ul>
<b>Arecanut</b>	<b>Fruiting</b>	<ul style="list-style-type: none"> <li>Due to clear sky and increase in temperature accelerate evaporation, provide irrigation to arecanut orchard at 6-8 days interval.</li> </ul>
<b>Chilli</b>	<b>Vegetative</b>	<ul style="list-style-type: none"> <li>There is possibility of incidence of leaf curl disease in chilli due to thrips, white flies and mites which results into curling of leaves and stunted growth of plant. For management spray 10% cow urine and Azadirachtin 10000 PPM @ 3 ml per liter of water, alternatively at an interval of 7 days after transplanting. Install blue sticky trap in field.</li> </ul>
<b>White onion</b>	<b>Nursery</b>	<ul style="list-style-type: none"> <li>For the nursery of white onion, carry out primary tillage operation. prepare raised bed of 3 m length x 1 m width x 15 cm height. Apply 5 kg well decomposed FYM, 35 gm urea, 100 gm single super phosphate and 25gm muriate of potash per sq.m at the time of sowing. Provide light irrigation after sowing.</li> </ul>
<p><b>This Agro Advisory Bulletin (AAB) is prepared and published with the consultation and recommendation of SMS committees of “Gramin Krishi Mausam Sewa (GKMS)” Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli. For more information contact nearby SAU research station or Agriculture officers of Agriculture Department, Maharashtra state.</b></p>		