



Ministry of Earth Science, India Meteorological Department is collaborated with
Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli,
Gramin Krishi Mousam Sewa Project,
College of Horticulture, Mulde
Agro Advisory Bulletin For The District Sindhudurg
(Period 18th to 22nd July, 2020)



Issue 56/2020

Date 17.07.2020

Duration 5 Days

Actual weather parameters recorded during last week
(Dated 10th to 16th July, 2020)

Weather Parameters

Forecasted weather parameters for forthcoming 5 day's
(Valid for 18th to 22nd July, 2020)

10/07	11/07	12/07	13/07	14/07	15/07	16/07	Date	18/07	19/07	20/07	21/07	22/07
26.6	67.0	168.0	12.8	19.8	52.4	57.6	Rainfall (mm)	46	31	30	15	8
29.5	31.4	29.0	27.0	27.0	26.0	26.0	Temp. Maximum (°C)	28	29	29	30	30
25.0	22.2	22.6	23.0	23.5	23.0	23.0	Temp. Minimum (°C)	24	25	25	25	24
Cloudy	Cloudy	Cloudy	Cloudy	Cloudy	Cloudy	Cloudy	AM	6	6	8	8	8
Cloudy	Cloudy	Cloudy	Cloudy	Cloudy	Cloudy	Cloudy	PM					
95	92	100	96	96	93	96	RH - I (%)	93	92	91	90	90
83	69	82	96	98	98	98	RH - II (%)	87	80	77	68	65
0.2	0.2	0.8	0.1	0.2	0.0	0.0	Wind Speed (km/hr.)	009	005	003	003	004
Calm	Calm	Calm	Calm	Calm	Calm	Calm	AM	South West	Southerly /South West	South	Westerly/ South West	Westerly/ South West
ENE	ENE	Calm	WNW	Calm	Calm	Calm	PM					

Rainfall (mm) in last year (2019)	Rainfall (mm) from 01/01/20 to till dated	Rainfall (mm) from 01/06/20 to till dated	Rainfall (mm) in last week
4591.0	2191.0	2187.8	404.2

General Weather condition in forthcoming 5 days:

In Sindhudurg district, the temperature is expected to drop and remain stable for the next five days. With moderate rains are expected in most places on July 18, 19 and 20 and light rains in most places on July 21st and 22nd. The weather is likely humid and completely cloudy. This forecast has been made by the Regional Meteorological Center, Mumbai. Warning: In Sindhudurg district on 18th July there is a possibility of heavy to very heavy rain in isolated places and heavy rain in isolated places on 19th and 20th July in isolated places.

Weather Prediction Based Agro-Advice

Name of Crop	Growth Stage	Advisory
Paddy	Tillering	<ul style="list-style-type: none"> There is a forecast of moderate rains in the forthcoming week. Application of second dose of fertilizers i.e. 870 gm of Urea per R area, if the crop has completed one month after transplanting. Before application of fertilizers, complete weeding operation and remove water. After application of fertilizers store water in the field. Regular supervision of the rice field should be carried out for any incidence of pests or diseases. Maintain water level of 5 cm by bunding rice field. Cleaning of bunds and proper drainage of water should be done. Use of Urea-DAP briquettes as per the recommendation of university is advocated. It is advocated to maintain proper level of water in the rice field. For control of blast disease in rice, need based application of Carbendazim 10 gms or copper oxychloride 25 gms per 10 litres of water is suggested.
Finger Millet	Transplanting to seedling	If remaining, complete the transplanting operations up to the end of this month. Transplanting of the Finger Millet should be carried out in drizzling rain or in cloudy weather, planting should be done across the slope of land also to increase production transplant 30 days old seedling by the spacing of 20 cm between rows and 15 cm between plant and two seedling per hill should be planted.
Groundnut	Sowing	Complete the sowing of kharif groundnut as early as possible.
Mango	New flush	If the application of fertilizer is completed in Mango, it is advocated to apply paclobutrazol during 15 th July to 31 st August for regular bearing in Mango for trees having age more than 10 years. It is applied as per the canopy of tree i.e. East-West of the canopy is 5 meters and North-South length of canopy is 3 meters, the average canopy will be 4 meters. Chemical (cultar) contains 23 per cent active ingredient of paclobutrazol. So it should be given @ 3 ml/meter canopy of the tree. For 4 meters the dose will be 12 ml/tree. Calculated quantity of paclobutrazol should be mixed in 3 to 6 litres of water as per the canopy of the tree. Apply paclobutrazol solution below the tree canopy i.e. by making 25 to 30 holes 15 cm deep inside the fertilizer ring. In each hole apply 100 ml solution. Weeding should be done before and after application of paclobutrazol. There should be sufficient moisture in the soil during application. Since the humidity will increase in the next five days in the district. Protect the new flushing in mango from the incidence of anthracnose and powdery mildew diseases may found on mango. To control spray Thiophenate methyl and Carbendazim 1% @ 1 gm per 1 liter of water after rains stop is advocated.
Cashew nut	-	If the incidence CSRB is noticed in cashew, remove the grubs with the help of 15 mm chisel and soak the stem in to the paste of fipronil 5% SC @ 2 ml per liter of water (5 liter of water is sufficient for one tree) is advocated. Cut the dead wood and diseased branches of cashew and apply 1% Bordeaux paste to the cut surface.
Vegetables	Vegetative	Proper support should be given to gourd crops. Provide proper drainage to the Bhendi, Gawar etc. crops.
Dairy Animals and Goat	Vaccination to Cows and Buffaloes against Black quarter and H.S. disease under the supervision of veterinary officer is advocated.	

This Agro Advisory Bulletin (AAB) is prepared and published with the consolation and recommendation of SMS committee of "Gramin Krishi Mausam Sewa (GKMS)", Regional Fruit Research Station, Vengurle and College of Horticulture, Mulde Dr. B.S.Konkan Krishi Vidyapeeth, Dapoli 416 520 (MS).

Dr. P. C. Haldavaneekar
Associate Dean and Nodal Officer
Agro-Meteorological Field Unit (AMFU),
Gramin Krishi Mousam Sewa,
College of Horticulture, Mulde
Tal. Kudal Dist. Sindhudurg
(02362-244231, 244272)

Dr. Y. C. Muthal
Technical Officer
Agro-Meteorological Field Unit (AMFU),
Gramin Krishi Mousam Sewa,
College of Horticulture, Mulde
(02362-244231)