

Wind Speed (km/hr):

Total cloud cover (Octa)

Date: 03/10/2017 (02358) 282387

Duration – 5 days



Agromet Advisory Service Bulletin for Raigad District n D.C. Konkon Kuishi Vidyonooth & Dooional India Mate -- **h**_- **i**)

(Issued jointly by IAAS, Dr. B.S. KonkanKrishiVidyapeeth, & Regional India Meteorological Department, Mumbai)									
Dr. Subhash Chavan,		Dr. D.N. Jagtap,		Prof. Viresh Chavan,					
Head,		Nodal Officer,		Technical Officer,					
Department of Agronomy		Department of Agronomy	Department of Agronomy						
9422431067		9403988143		9422065344					
Significant past weather for the preceding week (Period -29/09/2016 to 03/10/2017)		Weather forecast until 08.30 hrs of 08/10/2017							
		PARAMETERS							
			Day-1	Day-2	Day-3	Day – 4	Day – 5		
			04/10	05/10	06/10	07/10	08/10		
Rainfall (mm):		Rainfall (mm)	0	0	0	1	9		
Total Rainfall (mm):		Kamian (mm)	0	0	0	1	9		
Total Rainfall (mm) (last year)		Maximum temperature (⁰ C)	32	32	31	31	30		
Maximum temperature (⁰ C)		Minimum temperature (⁰ C)	26	26	26	26	26		
Minimum temperature (⁰ C)		Total cloud cover (Octa)	4	6	8	6	8		
Morning RH (%)		Relative Humidity Max. (%)	76	73	73	78	82		
Afternoon RH (%)		Relative Humidity Min. (%)	64	61	55	63	70		

Wind Speed (Km/hr)

Wind Direction

006

39

006

54

004

64

009

170

011

171

: Agro-met Advisory: It is possibility rainfall of 7th and 8th October, 2017. Sky will cloudy from 4th to 8th October, 2017. Wind velocity range from 4 to 11 kmph from 4th to 8th October, 2017.

Crop	Crop Stage	Agromet Advisory
Rice	Maturity as well as grain filling stage	Harvesting of early rice mature varieties.
Mango and Cashew	Plantation	 Weeding should be followed in Mango and Cashew garden with the help of sickle/grass cutting machine or spray Glyphosate weedicide @ 3-4 ml per lit water for effective control of weeds. <u>Note</u> : To avoid grazing of animal on weedicide spray in mango and cashew garden.
Coconut, Arecanut		 Spraying the FYM pits with 0.2 % Carbaryl for control of adults and grubs of rhinoceros beetle. To control red palm weevil affected coconut, fill the holes made by RPW with 10 per cent Carbaryl dust and sand mixture. Prepare a slanting hole with the help of screw drill about 1 m height from ground level on tree trunk and pour about 20 ml of 36 % Monocrotophos or 20% Chlorphyriphos with the help of plastic funnel in the hole and close the hole with the help of cement.
Vegetable	Fruiting	• Install methyl ugenol "Rakshak" tarp @ 4 nos. per ha in cucurbitaceous vegetable
crop	stage	crops, for effective control of fruit fly.
Goats/ Milch animals		• Provide clean and hygienic drinking water and nutritious fodder and concentrates to farm animal.
Suggestion		• Contact nearby SAU Scientists or State Agril. Dept. for detail control measures against incidence of pest and diseases under adverse weather conditions.