



## Agromet Advisory Service Bulletin for Thane District

(Issued jointly by IAAS, Dr. B.S. KonkanKrishiVidyapeeth, &amp; Regional India Meteorological Department, Mumbai)

**Dr. Subhash Chavan,**  
Head,  
Department of Agronomy  
9422431067

**Dr. D.N. Jagtap,**  
Nodal Officer,  
Department of Agronomy  
9403988143

**Prof. Viresh Chavan,**  
Technical Officer,  
Department of Agronomy  
9422065344

Significant past weather for the preceding week  
(Period –05/01/2018 to 09/01/2018)

|                                 |  |
|---------------------------------|--|
| Rainfall (mm):                  |  |
| Total Rainfall (mm):            |  |
| Total Rainfall (mm) (last year) |  |
| Maximum temperature (°C)        |  |
| Minimum temperature (°C)        |  |
| Morning RH (%)                  |  |
| Afternoon RH (%)                |  |
| Wind Speed (km/hr):             |  |
| Total cloud cover (Octa)        |  |

Weather forecast until 08.30 hrs of 14/01/2018

| PARAMETERS                 | Day-1 | Day-2 | Day-3 | Day - 4 | Day - 5 |
|----------------------------|-------|-------|-------|---------|---------|
|                            | 10/12 | 11/12 | 12/01 | 13/01   | 14/01   |
| Rainfall (mm)              | 0     | 0     | 0     | 0       | 0       |
| Maximum temperature (°C)   | 27    | 28    | 29    | 29      | 30      |
| Minimum temperature (°C)   | 18    | 20    | 20    | 21      | 21      |
| Total cloud cover (Octa)   | 0     | 0     | 1     | 0       | 0       |
| Relative Humidity Max. (%) | 81    | 79    | 86    | 84      | 85      |
| Relative Humidity Min. (%) | 50    | 50    | 51    | 50      | 50      |
| Wind Speed (Km/hr)         | 005   | 006   | 004   | 005     | 003     |
| Wind Direction             | 84    | 85    | 79    | 83      | 73      |

**: Agro-met Advisory: 218**Sky will be slightly cloudy on 12<sup>th</sup> January, 2017. Wind velocity range from 3 to 6 kmph from 10<sup>th</sup> to 14<sup>th</sup> January, 2017.

| Crop  | Crop Stage            | Agromet Advisory  |
|---|-----------------------|---|
| <b>Groundnut/<br/>Mustard</b>                         | Emergence Stage       | <ul style="list-style-type: none"> <li>For cultivation of rabi groundnut preliminary operation should be carried out. Deep ploughing and addition of 10 to 15 tonnes of FYM/Compost in groundnut field. Use of 125 to 150 kg seed for sowing. Use TG-26, TAG-24, TPG-41, Konkan Gaurav, Konkan Tapora any one variety for sowing.</li> <li>Weather is suitable for sowing of 'Mustard'. Sowing of seed of variety '<b>Pusa bold</b>' and '<b>Varuna</b>' at the spacing 45cm x 10 cm is advocated. Apply 1 kg urea and 3 kg SSP per R area before sowing of seed.</li> </ul>  |
| <b>Sapota</b>   | <b>Fruiting Stage</b> | <ul style="list-style-type: none"> <li>If unmaturing fruit drops of sapota are observed spray combined fungicides of (Metalxyl-M 8% + Mancozeb 64%) @ 0.2% i.e. 20 g per 10 lit. of water on whole plant canopy.</li> <li>It is a possibility for incidence of sapota seed borer, spray deltamethrin 2.8% E.C. @ 10 ml per 10 liter of water alongwith sticker. (The insecticide are not under label claim).</li> <li>Some parts in the district the sapota fruits are mature size stage, harvest the mature fruits before 10 hours in the morning and after 16 hours in the evening with the help on Atul Sapota harvester.</li> </ul>   |
| <b>Coconut,<br/>Areca nut</b>                         | --                    | <ul style="list-style-type: none"> <li>For control of adults and grubs of rhinoceros beetle. dust the FYM pits with 2 % Methyl parathion powder after every 2 month interval.</li> <li>To control red palm weevil affected coconut, prepare a slanting hole with the help of screw drill about 1 m height from ground level on tree trunk and pour about 20 ml of 36 % Monocrotophos or 20% Chlorphyriphos with the help of plastic funnel in the hole and close the hole with the help of cement.</li> </ul>   |
| <b>Vegetable<br/>crop and<br/>watermelon<br/>crop</b> | --                    | <ul style="list-style-type: none"> <li>For control of bacterial wilt of chilli, brinjal and tomato seedling, spray Copper Oxychloride @ 20 to 25 g/10 lit of water on seedbed after sowing of seeds. Also apply Carbaryl or Lindane powder on side of seedbed to control against ant.</li> <li>There is possibility for incidence of powdery mildew, spray Hexaconazole 5% EC 5 ml or Sulphur 80% WP @ 20 g per 10 lit. of waters.</li> <li>For cultivation of watermelon ploughing and cloud crushing of land and furrow be prepared at 4 m length 90 cm apart the both side of furrow prepared 30 x 30 x30 cm size pit, apply 1 to 1.5 kg FYM + 10 gm Carbaryl (10%) Powder, mix thoroughly with soil, sowing of 3 to 4 seeds at the depth of 2 to 2.5 cm, thinning after 15 days emergence of seedling keep only 2 seedlings per pit. The 50 kg N + 50 kg P<sub>2</sub>O<sub>5</sub> and 50 kg K<sub>2</sub>O per ha was required. The 1/3<sup>rd</sup> dose of N and full dose of P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O be applied at the time of sowing and remaining N be apply 1 and 2 months after sowing at equal quantity.</li> </ul> |
| <b>Goats/ Milch<br/>animals</b>                       | --                    | <ul style="list-style-type: none"> <li>Provide clean and hygienic drinking water and nutritious fodder and concentrates to farm animal.</li> <li>Protect the animals/poultry birds from low temperature by providing curtains and electric bulbs as per the need.</li> </ul>  |
| <b>Suggestion</b>                                     | --                    | <ul style="list-style-type: none"> <li>Contact nearby SAU Scientists or State Agril. Dept. for detail control measures against incidence of pest and diseases under adverse weather conditions.</li> </ul>  |