

**Agromet Advisory Service Bulletin for Ratnagiri District**  
(Issued jointly by IAAS, Dr. B.S. KonkankrishiVidyapeeth, & Regional India Meteorological Department, Mumbai)**Dr. Subhash Chavan,**  
Head,  
Department of Agronomy  
9422431067**Dr. Vijay More,**  
Nodal Officer,  
Department of Agronomy  
9422374001**Prof. Viresh Chavan,**  
Technical Officer,  
Department of Agronomy  
9422065344**Significant past weather for the preceding week**  
(Period –06/04/2018 to 10/04/2018)

Rainfall (mm):	0.0
Total Rainfall (mm):	0.0
Total Rainfall (mm) (last year)	3633.5
Maximum temperature (°C)	32.0-35.8
Minimum temperature (°C)	17.5-20.0
Morning RH (%)	87-94
Afternoon RH (%)	73-95
Wind Speed (km/hr):	3.9-4.7
Total cloud cover (Octa)	0-4

**Weather forecast until 08.30 hrs of 15/04/2018**

PARAMETERS	Day-1	Day-2	Day-3	Day - 4	Day - 5
	11/04	12/04	13/04	14/04	15/04
Rainfall (mm)	0	0	0	0	0
Maximum temperature (°C)	35	37	37	37	37
Minimum temperature (°C)	23	22	24	24	26
Total cloud cover (Octa)	3	1	0	2	3
Relative Humidity Max. (%)	69	77	74	65	79
Relative Humidity Min. (%)	33	26	27	27	30
Wind Speed (Km/hr)	005	007	008	006	006
Wind Direction	224	217	214	270	322

**: Agro-met Advisory:**Sky will be slightly cloudy from 11<sup>th</sup>, 12<sup>th</sup>, 14<sup>th</sup> and 15<sup>th</sup> April, 2018. Wind velocity range from 5 to 8 kmph from 11<sup>th</sup> to 15<sup>th</sup> April, 2018.

Crop	Crop Stage	Agromet Advisory
<b>Groundnut</b>	<b>Flowering</b>	<ul style="list-style-type: none"> <li>Due to increasing rate of evaporation irrigate the groundnut crop at an interval of 8-10 days after sowing. Rolling of empty drum 45-60 days after sowing for better penetration of pegs &amp; pod setting is advocated.</li> </ul>
<b>Pulses crops</b>	<b>Harvesting</b>	<ul style="list-style-type: none"> <li>Harvest mature pulse crop early in morning or late in evening and dry it for 3 to 4 days in sunlight. Stored dried grain in proper manner.</li> </ul>
<b>Mango and Cashew</b>	<b>Fruit Development</b> <b>Harvesting</b>	<ul style="list-style-type: none"> <li>Mango fruits mature size stage, harvest the mature fruits before 10 hours in the morning and after 16 hours in the evening with the help on Nutan mango harvester at 80 to 85% maturity. Keep the harvested fruits in shade to prevent spongy tissue disease and from heat. Give treatment with fungicide and ethylene for ripening as per the recommendations. Packing of mango fruits in C.F.B. (corrugated fiber box) developed by B. S. Konkankrishi Vidyapeeth, Dapoli is advised. Transport of harvested fruits should be done preferably during night hours. Don not apply any insecticides/fungicides 8 days before harvesting of mango fruits.</li> <li>There is possibility of incidence of Mango fruit fly, install 'Rakshak fruit fly traps' developed by University @ 4 traps per ha. area.</li> <li>If incidence of mango hopper if it is noticed on mango crop, spray Diamethoate 30 % EC @ of 10 ml or Lambda Cyhalothrin @5% E.C. @ of 6 ml per 10 liter of water</li> <li>It is necessary to protect fruit set of cashew from tea mosquito bug and thrips by spraying it with 50 % Profenophos 10 ml or 5% Lambda cyhalothrin @ 5 ml per 10 litres of water. (These insecticide are not under label claim)</li> <li>In some places the nuts ready for harvesting. Harvest the matured nuts and dry under sunlight.</li> <li>Prepare different processed product from cashew apple as per recommended by Dr. Balasaheb Sawant Konkankrishi Vidyapeeth, Dapoli.</li> </ul>
<b>Coconut, Arecanut</b>	--	<ul style="list-style-type: none"> <li>As there is increase in average temperature, irrigation should be given to coconut orchard at 5-6 days interval and for arecanut orchard at 4-5 days interval.</li> <li>For control of adults and grubs of rhinoceros beetle, dust 2 % Methyl parathion powder after every 2 month interval in FYM pits.</li> <li>To control red palm weevil affected coconut, prepare a slanting hole with the help of screw drill about 1 m height from ground level on tree trunk and pour about 20 ml of 36 % Monocrotophos or 20% Chlorphyriphos with the help of plastic funnel in the hole and close the hole with the help of cement.</li> </ul>
<b>Vegetable crop</b>	<b>Fruiting Stage</b>	<ul style="list-style-type: none"> <li>If incidence of hopper, aphids and thrips on vegetable crops viz., brinjal, tomato, cabbage, chilli, knol knol etc., is noticed spray Malathion @ 20 ml or Dimethoate @ 12 ml per 10 liter of water.</li> <li>Cucurbitaceous vegetables are in fruit bearing stage, install of Rakshak fruit fly traps developed by University @ 4 traps per ha area is advocated to control fruit fly in vegetables garden.</li> </ul>
<b>Goats/ Milch animals/ Poultry</b>	--	<ul style="list-style-type: none"> <li>Provide clean and hygienic drinking water and nutritious fodder and concentrates to farm animal.</li> <li>Provide 1 to 1.5 kg concentrate + 15 to 20 kg green fodder + 4 to 5 kg roughages per day for milch animals.</li> <li>Young goats provide 3 to 4 kg green fodder + 2 to 2.5 kg dry fodder + 300 g concentrate,.</li> <li>Protect the animals from afternoon heat. Keep good aeration in animal shed.</li> <li>For boiler poultry bird 1<sup>st</sup> three week provide boiler starter and 4 to 6 week old bird provide boiler finisher as per their daily requirement.</li> <li>The temperatures during day are increasing hence protect poultry birds from heat.</li> </ul>
<b>Suggestion</b>	--	<ul style="list-style-type: none"> <li>Contact nearby SAU Scientists or State Agril. Dept. for detail control measures against incidence of pest and diseases under adverse weather conditions.</li> </ul>