



Agromet Advisory Service Bulletin for Raigad District
(Issued jointly by IAAS, Dr. B.S. KonkanKrishiVidyapeeth, & Regional India Meteorological Department, Mumbai)

Dr. Subhash Chavan,
Head,
Department of Agronomy
9422431067

Dr. D.N. Jagtap,
Nodal Officer,
Department of Agronomy
9403988143

Prof. Viresh Chavan,
Technical Officer,
Department of Agronomy
9422065344

Significant past weather for the preceding week
(Period –08/01/2018 to 12/01/2018)

Rainfall (mm):	
Total Rainfall (mm):	
Total Rainfall (mm) (last year)	
Maximum temperature (°C)	
Minimum temperature (°C)	
Morning RH (%)	
Afternoon RH (%)	
Wind Speed (km/hr):	
Total cloud cover (Octa)	

Weather forecast until 08.30 hrs of 17/01/2018

PARAMETERS	Day-1	Day-2	Day-3	Day - 4	Day - 5
	13/01	14/01	15/01	16/01	17/01
Rainfall (mm)	0	0	0	0	0
Maximum temperature (°C)	34	34	33	32	32
Minimum temperature (°C)	22	21	21	20	19
Total cloud cover (Octa)	2	0	0	1	2
Relative Humidity Max. (%)	64	64	69	63	62
Relative Humidity Min. (%)	31	29	29	29	29
Wind Speed (Km/hr)	004	004	004	004	005
Wind Direction	110	60	36	77	129

: Agro-met Advisory:

Sky will be slightly cloudy on 13th, 16th and 17th January, 2017. Wind velocity range from 4 to 5 kmph from 13th to 17th January, 2017.

Crop	Crop Stage	Agromet Advisory
Summer Rice	Nursery	<ul style="list-style-type: none"> After 15 days of sowing of paddy nursery, apply second dose of nitrogenous fertilizer like urea @ 1 kg per R for paddy nursery.
Groundnut/ Mustard	Emergence Stage	<ul style="list-style-type: none"> For cultivation of <i>rabi</i> groundnut preliminary operation should be carried out. Deep ploughing and addition of 10 to 15 tonnes of FYM/Compost in groundnut field. Use of 125 to 150 kg seed for sowing. Use TG-26, TAG-24, TPG-41, Konkan Gaurav, Konkan Tapora any one variety for sowing. Weather is suitable for sowing of 'Mustard'. Sowing of seed of variety 'Pusa bold' and 'Varuna' at the spacing 45cm x 10 cm is advocated. Apply 1 kg urea and 3 kg SSP per R area before sowing of seed.
Mango and Cashew	Plantation	<ul style="list-style-type: none"> If incidence of mango hopper is noticed spray Imidachloprid 17.8% SL. @ of 3 ml per 10 liter of water for management of powdery mildew mix Hexaconazole 5% EC @ 5 ml or Sulphur 80% WP @ 20 gm per 10 lit of water. If incidence of Tea mosquito on new emerging flush of cashew is noticed, spray Lambda Cyhalothrin 5% EC (6 ml. in 10 lit. of water). (These insecticide are not under label claim)
Coconut, Arecanut	--	<ul style="list-style-type: none"> For control of adults and grubs of rhinoceros beetle. dust the FYM pits with 2 % Methyl parathion powder after every 2 month interval. To control red palm weevil affected coconut, prepare a slanting hole with the help of screw drill about 1 m height from ground level on tree trunk and pour about 20 ml of 36 % Monocrotophos or 20% Chlorphyriphos with the help of plastic funnel in the hole and close the hole with the help of cement.
Vegetable crop and watermelon crop	--	<ul style="list-style-type: none"> For control of bacterial wilt of chilli, brinjal and tomato seedling, spray Copper Oxychloride @ 20 to 25 g/10 lit of water on seedbed after sowing of seeds. Also apply Carbarly or Lindane powder on side of seedbed to control against ant. There is possibility for incidence of powdery mildew, spray Hexaconazole 5% EC 5 ml or Sulphur 80% WP @ 20 g per 10 lit. of waters. For cultivation of watermelon ploughing and cloud crushing of land and furrow be prepared at 4 m length 90 cm apart the both side of furrow prepared 30 x 30 x30 cm size pit, apply 1 to 1.5 kg FYM + 10 gm Carbaryl (10%) Powder, mix thoroughly with soil, sowing of 3 to 4 seeds at the depth of 2 to 2.5 cm, thinning after 15 days emergence of seedling keep only 2 seedlings per pit. The 50 kg N + 50 kg P₂O₅ and 50 kg K₂O per ha was required. The 1/3rd dose of N and full dose of P₂O₅ and K₂O be applied at the time of sowing and remaining N be apply 1 and 2 months after sowing at equal quantity.
Goats/ Milch animals	--	<ul style="list-style-type: none"> Provide clean and hygienic drinking water and nutritious fodder and concentrates to farm animal. Protect the animals/poultry birds from low temperature by providing curtains and electric bulbs as per the need.

Suggestion	--	<ul style="list-style-type: none">• Contact nearby SAU Scientists or State Agril. Dept. for detail control measures against incidence of pest and diseases under adverse weather conditions.
-------------------	----	--