



Agromet Advisory Service Bulletin for Raigad District

(Issued jointly by IAAS, Dr. B.S. KonkankrishiVidyapeeth, & Regional India Meteorological Department, Mumbai)

Dr. Subhash Chavan,
Head,
Department of Agronomy
9422431067

Dr. Vijay More,
Nodal Officer,
Department of Agronomy
9422374001

Prof. Viresh Chavan,
Technical Officer,
Department of Agronomy
9422065344

Significant past weather for the preceding week
(Period –09/03/2018 to 13/03/2018)

Rainfall (mm):	
Total Rainfall (mm):	
Total Rainfall (mm) (last year)	
Maximum temperature (^o C)	
Minimum temperature (^o C)	
Morning RH (%)	
Afternoon RH (%)	
Wind Speed (km/hr):	
Total cloud cover (Octa)	

Weather forecast until 08.30 hrs of 18/03/2018

PARAMETERS	Day-1	Day-2	Day-3	Day - 4	Day - 5
	14/03	15/03	16/03	17/03	18/03
Rainfall (mm)	0	0	3	3	0
Maximum temperature (^o C)	31	30	30	30	30
Minimum temperature (^o C)	22	22	22	22	22
Total cloud cover (Octa)	3	5	7	3	0
Relative Humidity Max. (%)	60	62	64	67	78
Relative Humidity Min. (%)	18	21	36	38	34
Wind Speed (Km/hr)	006	006	006	006	002
Wind Direction	61	301	85	81	83

: Agro-met Advisory:

It is possibility rainfall of 16th and 17th March, 2018. Sky will be cloudy from 14th to 17th March, 2018. Wind velocity range from 2 to 6 kmph from 14th to 18th March, 2018.

Crop	Crop Stage	Agromet Advisory
Summer Rice	Seedling	<ul style="list-style-type: none"> Apply 1st split dose of urea @ 55 kg/ha and maintained water level on rice field upto level 5 cm. before application of urea weeding should be carried out.
Pulses crops	Maturity stage	<ul style="list-style-type: none"> Harvest mature pulse crop early in morning or late in evening and dry it for 3 to 4 days in sunlight. Due to forecasted rainfall on 16th to 18th March, 2018, protect dried pulse crop from rain.
Mango and Cashew	Plantation	<ul style="list-style-type: none"> Due to temperature variation it is possibility for fruit drop of mango hence, apply irrigation for alphonso mango @ 150 to 200 lit. of water at fortnight interval to reduce fruit drop and increase the size of fruits. For increasing the production and quality improvement of fruits of mango, spraying of 1 % urea and 1 % Potassium nitrate at pea stage, marble stage and arecanut size stage is recommended. To avoid damage from increasing temperature as well as damage from fruit fly, wrap paper bags to marble to development size clear mango fruit. If incidence of mango hopper is noticed on mango crop, spray Diamethoate 30 % EC @ of 10 ml or Lambda Cyhalothrin @5% E.C. @ of 6 ml per 10 liter of water for management of Anthracnose mix Carbendazim @ 10 gm or Propineb @ 20 gm or Carbendazim 12% + Mancozeb 63% @ 10 g per of any one fungicides for10 lit of water. There is possibility of incidence of Mango fruit fly, install 'Rakshak fruit fly traps' developed by University @ 4 traps per ha. area. If incidence of Tea mosquito on new emerging flush of cashew is noticed, spray Lambda Cyhalothrin 5% EC (6 ml. in 10 lit. of water) also management of Anthracnose mix Carbendazim @ 10 gm or Propineb @ 20 gm or Carbendazim 12% + Mancozeb 63% @ 10 g per of any one fungicides for10 lit of water. (These insecticide are not under label claim)
Coconut, Arecanut	--	<ul style="list-style-type: none"> For control of adults and grubs of rhinoceros beetle, dust 2 % Methyl parathion powder after every 2 month interval in FYM pits. To control red palm weevil affected coconut, prepare a slanting hole with the help of screw drill about 1 m height from ground level on tree trunk and pour about 20 ml of 36 % Monocrotophos or 20% Chlorphyriphos with the help of plastic funnel in the hole and close the hole with the help of cement. Spray 1% Bordeaux mixture before rains or apply osety AL 0.3% (30 g/10 lit. water) + Urea Suphala 111 briquettes or Fosety AL 0.3% (30 g/10 lit. water) + Annapurana 76 briquettes per arecanut palm to avoid fungal immature fruit drop in coconut and Koleroga disease (immature nut drop) in arecanut.
Vegetable crop	--	<ul style="list-style-type: none"> If incidence of hopper, aphids and thrips on vegetable crops viz., brinjal, tomato, cabbage, chilli, knol knol etc., is noticed spray Malathion @ 20 ml or Dimethoate @ 12 ml per 10 liter of water. Cucurbitaceous vegetables are in fruit bearing stage, install of Rakshak fruit fly traps developed by University @ 4 traps per ha area is advocated to control fruit fly in vegetables garden.
Goats/ Milch animals/ Poultry	--	<ul style="list-style-type: none"> Due to forecasted rainfall on 16th to 18th March, 2018 protect stored feed and fodder from rain. Provide clean and hygienic drinking water and nutritious fodder and concentrates to farm animal. Planting of biennial and perennial forage crops for livestock like paragrass, hybrid napier, CO-3, yashwant etc. Provide 1 to 1.5 kg concentrate + 15 to 20 kg green fodder + 4 to 5 kg roughages per day for milch animals. Young goats provide 3 to 4 kg green fodder + 2 to 2.5 kg dry fodder + 300 g concentrate,. For boiler poultry bird 1st three week provide boiler starter and 4 to 6 week old bird provide boiler finisher as per their daily requirement.
Suggestion	--	<ul style="list-style-type: none"> Contact nearby SAU Scientists or State Agril. Dept. for detail control measures against incidence of pest and diseases under adverse weather conditions.