Date: 29/09/2017 (02358) 282387

Duration – 5 days



Agromet Advisory Service Bulletin for Thane District

(Issued jointly by IAAS, Dr. B.S. KonkanKrishiVidyapeeth, & Regional India Meteorological Department, Mumbai)

Dr. Subhash Chavan, Head, Department of Agronomy 9422431067

Dr. D.N. Jagtap,Nodal Officer, Department of Agronomy 9403988143

Prof. Viresh Chavan, Technical Officer, Department of Agronomy 9422065344

Significant past	weather for t	the preceding	g week
(Period)	-25/09/2017 t	o 29/09/2017)

Rainfall (mm):	
Total Rainfall (mm):	
Total Rainfall (mm) (last year)	
Maximum temperature (⁰ C)	
Minimum temperature (⁰ C)	
Morning RH (%)	
Afternoon RH (%)	
Wind Speed (km/hr):	
Total cloud cover (Octa)	

DADAMETEDO					
PARAMETERS	Day-1	Day-2	Day-3	Day - 4	Day - 5
	30/09	01/10	02/10	03/10	04/10
Rainfall (mm)	3	0	0	0	0
Maximum temperature (⁰ C)	35	36	36	36	36
Minimum temperature (⁰ C)	27	27	27	27	27
Total cloud cover (Octa)	6	4	3	6	6
Relative Humidity Max. (%)	86	83	85	79	71
Relative Humidity Min. (%)	62	58	56	53	50

006

135

007

130

007

126

008

130

006

121

Weather forecast until 08.30 hrs of 04/10/2017

: Agro-met Advisory: 218

Wind Speed (Km/hr)

Wind Direction

It is possibility rainfall of 30th September, 2017. Sky will be cloudy from 30th September 4th October, 2017. Wind velocity range from

Crop	Crop Stage	Agromet Advisory
Rice		• When intensity of rainfall very low apply 3 rd split dose of fertilizer to the late rice varieties
		at the time of flowering of rice crop @ 20 Kg N/ha.
		• There is a possibility for incidence of blue beetle, if it is noticed; spray Quinalphos 25%
		EC @ 2 lit. or or Triazophos 40% EC @ 625 ml or lambdacyhalothrin 5% EC @ 250 ml
		in 500 lit. of water per hector.
		• There is a possibility for incidence of fungal disease of blast on rice crop, if incidence is
		noticed; spray any one fungicides of Tricyclazole (75% WP) @ 10 gm per 10 lit of water
		or Isoprothiolane 40% @ 10 ml per 10 lit or water.
Sapota	Fruiting	• If unmature fruit drops of sapota are observed spray combined fungicides of (Metalxyl-M 8% +
	Stage	Mancozeb 64%) @ 0.2% i.e. 20 g per 10 lit. of water on whole plant canopy.
		• Some parts in the district the sapota fruits are mature size stage, harvest the mature fruits before 10
		hours in the morning and after 16 hours in the evening with the help on Atul Sapota harvester.
		Weeding should be followed in Sapota garden with the help of sickle/grass cutting machine or spray
		Glyphosate weedicide @ 3-4 ml per lit water for effective control of weeds.
		<u>Note</u> : To avoid grazing of animal on weedicide spray in mango and cashew garden.
Coconut,		• Spraying the FYM pits with 0.2 % Carbaryl for control of adults and grubs of rhinoceros beetle.
Arecanut		• To control red palm weevil affected coconut, fill the holes made by RPW with 10 per cent Carbaryl
		dust and sand mixture. Prepare a slanting hole with the help of screw drill about 1 m height from
		ground level on tree trunk and pour about 20 ml of 36 % Monocrotophos or 20% Chlorphyriphos
		with the help of plastic funnel in the hole and close the hole with the help of cement.
		• Apply osety AL 0.3% (30 g/10 lit. water) + Urea Suphala 111 briquettes or Fosety AL 0.3% (30 g/10 lit. water) + Annapurana 76 briquettes per arecanut palm to avoid fungal immature fruit drop in
		coconut and Koleroga disease (immature nut drop) in arecanut.
Vegetable	Fruiting	Install methyl ugenol "Rakshak" tarp @ 4 nos. per ha in cucurbitaceous vegetable crops, for
crop	stage	effective control of fruit fly.
Goats/		Provide clean and hygienic drinking water and nutritious fodder and concentrates to farm animal.
Milch		7,8
animals		
Suggestion		• Contact nearby SAU Scientists or State Agril. Dept. for detail control measures against incidence of
		pest and diseases under adverse weather conditions.