

**Agromet Advisory Service Bulletin for Raigad District**
(Issued jointly by IAAS, Dr. B.S. KonkanKrishiVidyapeeth, & Regional India Meteorological Department, Mumbai)**Dr. Subhash Chavan,**
Head,
Department of Agronomy
9422431067**Dr. D.N. Jagtap,**
Nodal Officer,
Department of Agronomy
9403988143**Prof. Viresh Chavan,**
Technical Officer,
Department of Agronomy
9422065344**Significant past weather for the preceding week**
(Period –27/10/2016 to 31/10/2017)

Rainfall (mm):	
Total Rainfall (mm):	
Total Rainfall (mm) (last year)	
Maximum temperature (⁰ C)	
Minimum temperature (⁰ C)	
0Morning RH (%)	
Afternoon RH (%)	
Wind Speed (km/hr):	
Total cloud cover (Octa)	

Weather forecast until 08.30 hrs of 05/11/2017

PARAMETERS	Day-1	Day-2	Day-3	Day - 4	Day - 5
	01/11	02/11	03/11	04/11	05/11
Rainfall (mm)	0	0	0	0	0
Maximum temperature (⁰ C)	37	37	37	36	35
Minimum temperature (⁰ C)	22	23	23	22	23
Total cloud cover (Octa)	0	2	5	0	0
Relative Humidity Max. (%)	61	61	63	65	62
Relative Humidity Min. (%)	31	32	32	35	33
Wind Speed (Km/hr)	006	006	006	006	008
Wind Direction	81	77	76	70	74

: Agro-met Advisory:Sky will slightly cloudy on 2nd and 3rd November, 2017. Wind velocity range from 6 to 8 kmph from 01st to 5th November, 2017.

Crop	Crop Stage	Agromet Advisory
Rice	Maturity stage	<ul style="list-style-type: none"> Harvesting of rice crop in morning hours and fallowed immediate threshing of rice, the threshed grains should be dry in sun for 3 to 4 days.
Mango and Cashew	Plantation	<ul style="list-style-type: none"> There is a possibility for incidence of mango hopper, if it is noticed spray Deltamethrin 2.8% E.C. @ of 9 ml per 10 liter of water alongwith sticker. There is a possibility for incidence of Tea mosquito bug on cashew, if it is noticed spray Lambda Cyhalothrin 5% E.C. @ 6 ml per 10 liter of water alongwith sticker (The insecticide are not under label claim).
Coconut, Arecanut	--	<ul style="list-style-type: none"> For controlling of bud rot of coconut palm, clean infected bud and drenching with 1% Bordo mixture or copper-oxy-chloride @ 25 g per 10 lit of water. Spraying the FYM pits with 0.2 % Carbaryl for control of adults and grubs of rhinoceros beetle. To control red palm weevil affected coconut, fill the holes made by RPW with 10 per cent Carbaryl dust and sand mixture. Prepare a slanting hole with the help of screw drill about 1 m height from ground level on tree trunk and pour about 20 ml of 36 % Monocrotophos or 20% Chlorphyriphos with the help of plastic funnel in the hole and close the hole with the help of cement.
Vegetable crop	Fruiting stage	<ul style="list-style-type: none"> Install methyl ugenol "Rakshak" tarp @ 4 nos. per ha in cucurbitaceous vegetable crops, for effective control of fruit fly. Prepare nursery of tomato, brinjal and chilly seedling for arriving <i>rabi</i> season.
Goats/ Milch animals	--	<ul style="list-style-type: none"> Provide clean and hygienic drinking water and nutritious fodder and concentrates to farm animal. Protect the poultry birds from low temperature by providing electric bulbs as per the need.
Suggestion	--	<ul style="list-style-type: none"> Contact nearby SAU Scientists or State Agril. Dept. for detail control measures against incidence of pest and diseases under adverse weather conditions.